



Phylogeography of the Japanese Giant Flying Squirrel, *Petaurista leucogenys*, Based on Mitochondrial DNA Control Region Sequences

著者 (英)	Oshida Tatsuo, Ikeda Koichi, Yamada Kazuhiko, Masuda Ryuichi
journal or publication title	Zoological science
volume	18
number	1
page range	107-114
year	2001
URL	http://id.nii.ac.jp/1588/00004214/

doi: info:doi/10.2108/zsj.18.107

Phylogeography of the Japanese Giant Flying Squirrel, *Petaurista leucogenys*, Based on Mitochondrial DNA Control Region Sequences

Tatsuo Oshida^{1†}, Koichi Ikeda², Kazuhiko Yamada³
and Ryuichi Masuda^{1*}

¹Chromosome Research Unit, Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan

²Fukuoka Prefecture Forest Research and Extension Center, Kurume 839-0827, Japan

³Laboratory of Cytogenetics, Division of Biosciences, Graduate School of Environmental Earth Science, Hokkaido University, Sapporo 060-0810, Japan

ABSTRACT—To investigate genetic diversity among populations of the Japanese giant flying squirrel *Petaurista leucogenys*, the mitochondrial DNA control region sequences (1,052–1,054 bases) were determined in 37 specimens from 17 localities on the Honshu, Shikoku, and Kyushu Islands of Japan. Of the 37 animals examined, 24 haplotypes were identified. All haplotypes from Kyushu consisted of 1,052 bases, whereas those from Honshu and Shikoku consisted of 1,054 bases including two insertions, except for three haplotypes (which had 1,052 or 1,053 bases). Phylogenetic relationships reconstructed using neighbor-joining and maximum parsimony methods indicated that *P. leucogenys* is essentially separated into three major lineages: Group A consisting of a single haplotype from Kyushu, Group B consisting of some haplotypes from Kyushu and one haplotype from Honshu, and Group C consisting mostly of haplotypes from Honshu and Shikoku. Animals with the Kyushu haplotypes were split into two lineages (Groups A and B), suggesting that Group A diverged at an earlier point from the other groups. Genetic distances in Group C were not related to geographic distances between sampling localities, indicating that ancestral populations of this group recently expanded their distribution in a short time, possibly after the last glacial stage.

INTRODUCTION

The Japanese giant flying squirrel *Petaurista leucogenys* is a mammalian species indigenous to the Kyushu, Shikoku, and Honshu Islands of Japan (Corbet and Hill, 1991; Nowak, 1991). Although Corbet and Hill (1980) explained that this species is distributed throughout Japan and central China, they more recently treated the central Chinese population as a distinct species *Petaurista xanthotis* (Corbet and Hill, 1991; 1992). Based on characteristics of their pelage, Imaizumi (1960) classified *P. leucogenys* into three subspecies: *leucogenys*, *nikkonis*, and *oreas*. At present, however, the validity of his classification is uncertain, as there are variations of pelage in *P. leucogenys*.

The ecology of *P. leucogenys* has been studied in detail

(Baba *et al.*, 1982; Ando and Imaizumi, 1982; Ando and Shiraishi, 1983; Kawamichi, 1997a; 1997b; 1998), and cytogenetic information on this species has also been reported by Oshida and Obara (1991; 1993) and Oshida and Yoshida (1999a; 1999b). However, little information about the phylogeography of *P. leucogenys* has been known heretofore. Oshida and Obara (1993) reported the variation of constitutive heterochromatin of chromosomes in *P. leucogenys*, but they did not find any geographically specific features.

In the present study, in order to study the phylogeography and subspecies classification of *P. leucogenys*, we analyzed the mitochondrial DNA (mtDNA) control region sequences. Since the control region contains variable blocks which evolve about 4–5 times faster than the other regions of mtDNA molecules (Greenberg *et al.*, 1983; Horai and Hayasaka, 1990; Brown *et al.*, 1993), this region is a very valuable molecular marker for investigating relationships among closely related species or conspecific populations (e.g., Baker *et al.*, 1993; Arctander *et al.*, 1996; Nagata *et al.*, 1998; 1999; Barratt *et al.*, 1999; Kurose *et al.*, 1999; Matsushashi *et al.*, 1999). Based

* Corresponding author: Tel. +81-11-706-3541;
FAX. +81-11-736-6304.

E-mail: masudary@ees.hokudai.ac.jp

† Present address: Laboratory of Molecular Ecology, Department of Biology, Tunghai University, Taichung Taiwan 407, R. O. C.

on the control region data, we here discuss phylogeographic relationships within Japanese populations of *Petaurista leucogenys*.

MATERIALS AND METHODS

Animals

A profile of the Japanese giant flying squirrel *Petaurista leucogenys* examined in the present study is shown in Table 1. Thirty seven specimens of *P. leucogenys* were collected from 17 localities in Japan (Fig. 1). A female red giant flying squirrel *Petaurista petaurista melanotus* (PPM) imported from Hong-Kong to Japan in 1990 was used as an out-group.

DNA preparation and sequencing

Total DNAs were extracted from muscle or liver tissues using the phenol/proteinase K/sodium dodecyl sulfate method of Sambrook *et al.* (1989). The whole control region was amplified using polymerase chain reaction (PCR), with a set of newly designed primers: L15933 5'-CTCTGGTCTTGTAACCAAAAATG-3' and H637 5'-AGGACC-

AAACCTTTGTGTTTATG-3'. Primer names correspond to the light (L) or heavy (H) strand and the 3'end-position of the primers in the human mtDNA sequences (Anderson *et al.*, 1981). The reaction mixture of 50 µl contained 100 ng of genomic DNA, 25 picomoles of each primer, 200 µM dNTPs, 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl₂, and 2.5 units of *rTaq* DNA polymerase (Takara). Amplification was carried out for 35 cycles as follows: 94°C for 1 min, 50°C for 1 min, and 72°C for 2 min, and then the extension reaction was performed at 72°C for 10 min. PCR products were purified with the Qia-quick PCR purification kit (QIAGEN) and directly sequenced using an automated DNA sequencer (SQ5500L, Hitachi). For sequencing, two PCR primers and another new primer (5'-CCTAATGGATATCCCCTTCCAACG-3') were used.

Phylogenetic analysis

All sequences were aligned using the computer software GeneWorks (Intelligenetics). The phylogenetic tree was constructed via the neighbor-joining (NJ) method (Saitou and Nei, 1987) in Clustal W (Thompson *et al.*, 1994) and via the maximum parsimony (MP) method using a heuristic search algorithm with the 50% majority-rule consensus in PAUP (Swofford, 1993). In the NJ tree, the numbers of

Table 1. Specimen profiles of *Petaurista leucogenys* examined in the present study

Sample name	Sex#	Sampling locality (Supplier)	No.** of locality	Haplotype	Accession No. of sequence***
AM1*	M	Sannohe-gun, Aomori Pref.	1	H6	AB043805
AM2*	F	Sannohe-gun, Aomori Pref.	1	H5	AB043804
IT1	M	Koromogawa, Iwate Pref. (Morioka Zoo)	2	H14	AB043813
IT2	F	Koromogawa, Iwate Pref. (Morioka Zoo)	2	H12	AB043811
NN1*	M	Shiga-Height, Nagano Pref.	3	H11	AB043810
NN2*	M	Shiga-Height, Nagano Pref.	3	H4	AB043803
NN3*	M	Shiga-Height, Nagano Pref.	3	H13	AB043812
TY1*	F	Nakanikawa-gun, Toyama Pref.	4	H13	AB043812
TY2*	M	Nakanikawa-gun, Toyama Pref.	4	H4	AB043803
YN1*	M	Nirazaki, Yamanashi Pref.	5	H4	AB043803
YN2*	F	Nirazaki, Yamanashi Pref.	5	H10	AB043809
TG1	M	Nikko, Tochigi Pref. (Tochigi Prefectural Museum)	6	H7	AB043806
TG2	M	Shioya, Tochigi Pref. (Tochigi Prefectural Museum)	7	H8	AB043807
KN1	M	Hakone, Kanagawa Pref. (Kanagawa Prefecture Natural Environment Conservation Center)	8	H15	AB043814
KN2	F	Aikawa, Kanagawa Pref. (Kanagawa Prefecture Natural Environment Conservation Center)	9	H16	AB043815
WK1*	M	Hashimoto, Wakayama Pref.	10	H9	AB043808
WK2*	F	Hashimoto, Wakayama Pref.	10	H4	AB043803
GF1*	M	Kamioka, Gifu Pref.	11	H4	AB043803
GF2*	F	Kamioka, Gifu Pref.	11	H5	AB043804
KT1	M	Kyoto, Kyoto Pref. (Mr. M. Kishioki)	12	H2	AB043801
HS1	F	Hiroshima Pref. (Asa Zoological Park)	13	H1	AB043800
KG1*	M	Takamatsu, Kagawa, Pref	14	H4	AB043803
EH1	?	Omgo, Ehime Pref. (Omogo Mountain Museum)	15	H3	AB043802
FO1	F	Hirokawa, Fukuoka Pref.	16	K1	AB043792
FO2	M	Hirokawa, Fukuoka Pref.	16	K1	AB043792
FO3	M	Hirokawa, Fukuoka Pref.	16	K1	AB043792
FO4	F	Joyo, Fukuoka Pref.	17	K5	AB043796
FO6	F	Joyo, Fukuoka Pref.	17	K1	AB043792
FO7	F	Joyo, Fukuoka Pref.	17	K3	AB043794
FO8	F	Joyo, Fukuoka Pref.	17	K8	AB043799
FO9	M	Joyo, Fukuoka Pref.	17	K2	AB043793
FO10	M	Joyo, Fukuoka Pref.	17	K4	AB043795
FO11	M	Joyo, Fukuoka Pref.	17	K2	AB043793
FO15	M	Joyo, Fukuoka Pref.	17	K5	AB043796
FO16	M	Joyo, Fukuoka Pref.	17	K1	AB043792
FO17	M	Joyo, Fukuoka Pref.	17	K6	AB043797
FO18	F	Joyo, Fukuoka Pref.	17	K7	AB043798

* Specimens obtained commercially from pet stores in Japan: Takita Store, Sannohe-gun and Saitama Sougou Pet, Koshigaya.

** Locality Nos. correspond to those in Fig.1.

*** Sequence data will appear in the DDBJ nucleotide sequence databases with these accession numbers.

M, male; F, female.




Fig. 1. Sampling localities in the present study and fossil localities of *Petaurista leucogenys*. Closed circles, sampling localities; open square, Middle Pleistocene fossil localities; closed squares, Late Pleistocene fossil localities. Sampling locality numbers correspond to those of Table 1 and Fig. 2. Arrow indicates the presumed migration route of *P. leucogenys* from southern China to Japan.

nucleotide substitutions per site were estimated for multiple substitutions by Kimura's (1980) two-parameter method. Using sequences without gap-sites, the MP tree was produced by unweighted parsimony. To assess the branching confidence, bootstrap values (Felsenstein, 1985) were derived from 1,000 replications of the NJ method and 100 replications of the MP tree.

RESULTS

Sequence Divergence of mtDNA Control Region

All mtDNA control regions (1,052–1,054 bases) of *P. leucogenys* from 17 localities in Japan were successfully sequenced. All sequences of animals from Kyushu had 1,052 bases, while those of populations from Honshu and Shikoku had 1,054 bases with insertions except for three specimens: EH1 (1,052 bases), KT1 (1,052 bases), and HS1 (1,053 bases) (Table 2). Of all sequences obtained, 145 sites were variable: transitions at 117, transversions at 14 sites, and both transitions and transversions at 12 sites (Table 2). In the 37 animals, 24 haplotypes were identified (Tables 1 and 2). The sequence divergence among haplotypes was 0.7–4.8% (Table 3).

The control region of *P. leucogenys* was divided into three domains: two variable domains (the 5' and 3' ends) and one conserved central domain (Table 2). The 5' end domain was more variable than the 3' end domain and contained one gap-site (site number 281) in all haplotypes from Kyushu, in H1 and H2 from Honshu, and in H3 from Shikoku. The 3' end domain contained an additional gap-site (site number 1,052)

in all haplotypes from Kyushu, in H2 from Honshu, and in H3 from Shikoku. The sequence of *P. petaurista* used as an out-group has 1,051 bases (accession number in DDBJ: AB043816).

Molecular phylogeny based on the mtDNA control region

Phylogenetic relationships reconstructed via NJ and MP methods were similar to each other. In the NJ tree, the Japanese population of *P. leucogenys* was separated into three major lineages: Group A consisting of K1, Group B consisting of K2, K3, K4, K5, K6, K7, K8, and H1 (95% bootstrap value), and Group C consisting of H4, H5, H6, H7, H8, H9, H10, H11, H12, H13, H14, and H15 (62% bootstrap value) (Fig. 2a). In the MP analysis, only one most-parsimonious phylogenetic tree was obtained by unweighted parsimony, and it had a consistency index of 0.668. The three major groups were also recognized in the MP tree: Group A (K1), Group B consisting of K2, K3, K4, K5, K6, K7, K8, and H1 (88% bootstrap value), and Group C consisting of H4, H5, H6, H7, H8, H9, H10, H11, H12, H13, H14, and H15 (57% bootstrap value) (Fig. 2b). In both trees, H1 from Honshu was clustered with the Kyushu population consisting of K2, K3, K4, K5, K6, K7, and K8 with high bootstrap values (95% in the NJ tree and 88% in the MP tree), and H2, H3, and H16 were not included in the major three lineages. In addition, in both trees, genetic differences between haplotypes of Group C did not correspond to geographic distances. The sequences of Groups A and B had

1,052 bases and those of Group C had 1,054 bases, although there was an exception (H1).

DISCUSSION

Characterization of the mtDNA control region in *Petaurista leucogenys*

In vertebrates, it has been reported that the control region consists of two variable domains (5' and 3' ends) and a conserved central domain (Brown *et al.*, 1986; Southern *et al.*, 1988; Saccone *et al.*, 1991). In the same way, the control region of *P. leucogenys* examined in the present study was also divided into three domains (Table 2). In particular, the 5' end domain was more variable than the 3' end domain and contained one gap-site (site number 281) in all haplotypes from Kyushu, in H1 and H2 from Honshu, and in H3 from Shikoku. Another gap-site (site number 1,052) was recognized in the 3' end domain in all haplotypes from Kyushu and in H2 and H3. The two gap-sites were specific to the control region of the Kyushu population. In addition, repetitive sequences, which were reported in the control region of some mammals (e.g., Hoelzel *et al.*, 1994; Nagata *et al.*, 1998; Kurose *et al.*, 1999; Matsuhashi *et al.*, 1999), were not found in *P. leucogenys* and *P. petaurista*.

Phylogeography of *Petaurista leucogenys*

According to pelage characteristics, Imaizumi (1960) classified *P. leucogenys* into three subspecies (*leucogenys*, *nikkonis*, and *oreas*), and demonstrated that *P. l. leucogenys* is distributed throughout the Kyushu and Shikoku islands, that *P. l. nikkonis* occurs in the eastern part of the Honshu Island, and that *P. l. oreas* occurs in the western part of the Honshu Island. However, in the present study, phylogenetic relationships among haplotypes did not correspond to subspecies classification and distribution.

Despite of the small number of specimens collected (Fig. 1), the Kyushu population was divided into two lineages: K1 lineage (Group A) and another lineage consisting of K2, K3, K4, K5, K6, K7, and K8 in Group B (Fig. 2). In the NJ and MP trees, K1 was likely to have been isolated from the other haplotypes by the first dichotomy, although the bootstrap values were not so high (< 50% in NJ tree; 59% in MP tree), suggesting that K1 diverged from the other haplotypes at an earlier point. On the other hand, although H1 was closely related to K2, K3, K4, K5, K6, K7, and K8 in both trees, the phylogenetic positions of H2 and H3 in the NJ tree were different from those in the MP tree (Fig. 2). The three haplotypes (H1, H2, and H3) shared the two gap-sites specific to the Kyushu population: site number 281 in H1, H2, and H3 and site number 1,052 in H2 and H3 (Table 2). This indicated that H2 and H3 are more closely related to K2, K3, K4, K5, K6, K7, and K8 or that they are intermediate haplotypes between the Kyushu population and Honshu/Shikoku population. The phylogenetic position of H16 that was not clustered with the three major lineages was not clear in the present study.

In terms of the Honshu/Shikoku population in the NJ and

MP trees, it is noteworthy that genetic distances between haplotypes do not always correspond to geographic distances of sampling localities, and exhibit low bootstrap values (Fig. 2). Moreover, six specimens from different localities (Gifu, Kagawa, Nagano, Toyama, Wakayama, and Yamanashi) shared the H4 haplotype, and two specimens from Aomori and Gifu had the H5 haplotype (Fig. 1 and Table 1). In the Honshu/Shikoku population, except for H1, H2, H3, and H16, haplotype H4 seemed to be most common.

Based on some fossil records, Kawamura (1988) reported that *P. leucogenys* had presumably migrated from southern China to Japan through the land bridge (Fig. 1) which was formed around the area of the present East China Sea in the Early Middle Pleistocene. It was not known that *Petaurista* had existed in the Korean Peninsula at that time. Accepting Kawamura's hypothesis (1988), inevitably, the first place where *P. leucogenys* had migrated from southern China could have been the Kyushu Island in Japan, and then it could have extended its distribution toward the Honshu and Shikoku Islands (see Fig. 1). Fossils of *Petaurista* before the Holocene period are very rare in the Japanese islands. The Middle Pleistocene fossils of *P. leucogenys* were found in two localities of Japan (Hasegawa, 1966; 1972; Kowalski and Hasegawa, 1976; Kawamura, 1988) which are very close to each other as shown in Fig. 1. Moreover, the Late Pleistocene fossils of *P. leucogenys* were also recognized in six localities of Japan (Shikama, 1949; Hasegawa, 1966; Kowalski and Hasegawa, 1976; Kawamura, 1980; 1981; 1982; 1988; Kawamura and Sotsuka, 1984; Kawamura *et al.*, 1986), as shown in Fig. 1. Judging from these fossil records, by the Middle or Late Pleistocene, *P. leucogenys* could already have been distributed in the Honshu and Shikoku Islands. *Petaurista leucogenys* is an arboreal animal and inhabits the temperate forests (Nowak, 1991). Therefore, during glacial stages in the Pleistocene, the habitats of this animal may have been reduced due to the southward shifting of temperate forests in Japan. The results of the present study show that the genetic distances in the Honshu/Shikoku population were not related to geographic distances of sampling localities. This suggests that *P. leucogenys* rapidly extended its distribution in a short time during the northward expansion of temperate forests in Japan after the last glacial stage of the Pleistocene.

Although the evolutionary rate of humans may not always correspond to that of the giant flying squirrel because of the differences of generation time between humans and giant flying squirrels, applying the evolutionary rate (approximately 8.4% per million years, Myr) of the human control region reported by Vigilant *et al.* (1989), the divergence times between Groups A and B, between Groups A and C, and between Groups B and C were estimated to be approximately 0.4–0.5, 0.4–0.5, and 0.4–1.0 Myrs ago, respectively. On the other hand, the divergence times in Group B and in Group C were approximately 0.1–0.3 and 0.1–0.5 Myrs ago, respectively. Accordingly, the divergences among haplotypes of *P. leucogenys* may have occurred rapidly from the Middle to Late Pleistocene.

ACKNOWLEDGMENTS

We would like to thank Dr. T. Kawamichi and Dr. S. Uni (Osaka City University), Dr. M. Baba (Kitakyushu Museum of Natural History), Mr. H. Kuribayashi, Dr. C. Katoh, and Dr. M. Itoh (Kanagawa Prefecture Natural Environment Conservation Center), Mr. T. Tsujimoto (Morioka Zoo), Mr. T. Okayama (Omogo Mountain Museum), Dr. S. Yachimori (Tochigi Prefectural Museum), Asa Zoological Park, Mr. M. Kishioki, and Ms. M. Shigeta, for supplying specimens. This study was partly supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Science, Sports, and Culture of Japan.

REFERENCES

- Anderson S, Bankier AT, Barrell BG, De Bruijn MHL, Coulson AR, Drouin J, Eperon IC, Nierlich DP, Roe BA, Sanger F, Schreier PH, Smith AJ H, Staden R, Young IG (1981) Sequence and organization of the human mitochondrial genome. *Nature* 290: 457–465
- Ando M, Imaizumi Y (1982) Habitat utilization of the white-cheeked giant flying squirrel *Petaurista leucogenys* in a small shrine grove. *J Mamm Soc Jpn* 9: 70–81 (in Japanese with English abstract)
- Ando M, Shiraishi S (1983) The nest and nest-building behavior of the Japanese giant flying squirrel, *Petaurista leucogenys*. *Sci Bull Fac Agr Kyushu Univ* 38: 59–69 (in Japanese with English abstract)
- Arctander P, Kat PW, Aman RA, Siegmund HR (1996) Extreme genetic differences among populations of *Gazella granti*, Grant gazelle, in Kenya. *Heredity* 76: 465–475
- Baba M, Doi T, Ono Y (1982) Home range utilization and nocturnal activity of the giant flying squirrel, *Petaurista leucogenys*. *Jpn J Ecol* 32: 189–198
- Baker AJ, Piersma T, Rosenmeier L, Weinrich MT, Anernethy RB, Calambokidis J, Lien J, Lamberstein RH, Ramirez JU, Vasquez O, Clapham PJ, Alling A, O'Brien SJ, Palumbi SR (1993) Abundant mitochondrial DNA variation and world-wide population structure in humpback whales. *Proc Natl Acad Sci USA* 90: 8239–8243
- Barratt EM, Gurnell J, Malarky G, Deaville R, Bruford MW (1999) Genetic structure of fragmented populations of red squirrel (*Sciurus vulgaris*) in the UK. *Mol Ecol* 8: S55–S63
- Brown GG, Gadaleta G, Pepe G, Saccone C, Sbsia E (1986) Structural conservation and variation in the D loop-containing region of vertebrate mitochondrial DNA. *J Mol Biol* 192: 503–511
- Brown JR, Beckenbach AT, Smith MJ (1993) Intraspecific DNA sequence variation of the mitochondrial control region of white sturgeon (*Acipenser transmontanus*). *Mol Biol Evol* 10: 326–341
- Corbet GB, Hill JE (1980) *A World List of Mammalian Species*. 1st ed, Oxford University Press, Oxford, pp 136–137
- Corbet GB, Hill JE (1991) *A World List of Mammalian Species*. 3rd ed, Oxford University Press, Oxford, pp 144–145
- Corbet GB, Hill JE (1992) *The Mammals of the Indomalayan Region: A Systematic Review*. Oxford University Press, Oxford, pp 308–313
- Felsenstein J (1985) Confidence limits on phylogenies: An approach using the bootstrap. *Evolution* 39: 783–791
- Greenberg BD, Newbold JE, Sugino A (1983) Intraspecific nucleotide sequence variability surrounding the origin of replication in human mitochondrial DNA. *Gene* 21: 33–49
- Hasegawa Y (1966) Quaternary smaller mammalian fauna from Japan. *Fossils* 11: 31–40 (in Japanese)
- Hasegawa Y (1972) The NAUMANN'S elephant, *palaeoloxodon naumanni* (MAKIYAMA) from the Late Pleistocene of Shakagahara, Shodoshima Is. in Seto Inland Sea, Japan. *Bull Natl Sci Mus* 15: 513–591
- Hoelzel AR, Lopez JV, Dover GA, O'Brien SJ (1994) Rapid evolution of a heteroplasmic repetitive sequence in the mitochondrial DNA control region of carnivores. *J Mol Evol* 39: 191–199
- Horai S, Hayasaka K (1990) Intraspecific nucleotide sequence differences in the major noncoding region of human mitochondrial DNA. *Am J Hum Genet* 46: 828–842
- Imaizumi Y (1960) *Colored Illustration of the Mammals of Japan*. Hoikusha, Osaka (in Japanese)
- Kawamichi T (1997a) Seasonal changes in the diet of Japanese giant flying squirrel in relation to reproduction. *J Mamm* 78: 204–212
- Kawamichi T (1997b) The age of sexual maturity in Japanese giant flying squirrel, *Petaurista leucogenys*. *Mammal Study* 22: 81–87
- Kawamichi T (1998) Seasonal change of testis size in Japanese giant flying squirrel *Petaurista leucogenys*. *Mammal Study* 23: 79–82
- Kawamura Y (1980) Mammalian remains of the Pre-Jomon Period from Taishaku-Kannondo Cave Site (Part 1). Mammalian remains obtained by the excavation of 1975. *Ann Bull Hiroshima Univ Taishaku-kyo Sites Res Centre* 3: 61–74 (in Japanese)
- Kawamura Y (1981) Mammalian remains of the Pre-Jomon Period from Taishaku-Kannondo Cave Site (Part 2). Mammalian remains obtained by the excavation of 1976. *Ann Bull Hiroshima Univ Taishaku-kyo Sites Res Centre* 4: 67–88 (in Japanese)
- Kawamura Y (1982) Mammalian remains of the Pre-Jomon Period from Taishaku-Kannondo Cave Site (Part 3). Mammalian remains obtained by the excavation of 1978. *Ann Bull Hiroshima Univ Taishaku-kyo Sites Res Centre* 5: 57–70 (in Japanese)
- Kawamura Y (1988) Quaternary rodent faunas in the Japanese Islands (Part 1). *Mem Fac Sci, Kyoto Univ, Ser Geol Min* 53: 31–348
- Kawamura Y, Sotsuka T (1984) Preliminary report on the Quaternary mammalian remains from several caves on the Hiraodai Plateau, Fukuoka Prefecture, Northern Kyushu, Japan. *Bull Kitakyushu Mus Nat Hist* 5: 163–188 (in Japanese with English abstract)
- Kawamura Y, Yamada Y, Ando Y (1986) Late Pleistocene micro-mammals from Taishaku-Kannondo Cave Site (first preliminary report). *Ann Bull Hiroshima Univ Taishaku-kyo Sites Res Centre* 9: 67–85 (in Japanese)
- Kimura M (1980) A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* 16: 111–120
- Kowalski K, Hasegawa Y (1976) Quaternary rodents from Japan. *Bull Nat Sci Mus Ser C* 2: 31–66
- Kurose N, Masuda R, Yoshida MC (1999) Phylogeographic variation in two Mustelinae, the least weasel *Mustela nivalis* and the ermine *M. erminea* of Japan, based on mitochondrial DNA control region sequences. *Zool Sci* 16: 971–977
- Matsuhashi T, Masuda R, Mano T, Yoshida MC (1999) Microevolution of the mitochondrial DNA control region in the Japanese brown bear (*Ursus arctos*) population. *Mol Biol Evol* 16: 676–684
- Nagata J, Masuda R, Kaji K, Kaneko M, Yoshida MC (1998) Genetic variation and population structure of the Japanese sika deer (*Cervus nippon*) in Hokkaido island, based on mitochondrial D-loop sequences. *Mol Ecol* 7: 871–877
- Nagata J, Masuda R, Tamate HB, Hamasaki S, Ochiai K, Asada M, Tatsuzawa S, Suda K, Tado H, Yoshida MC (1999) Two genetically distinct lineages of the sika deer, *Cervus nippon*, in Japanese islands: comparison of mitochondrial D-loop region sequences. *Mol Phylogenet Evol* 13: 511–519
- Nowak RM (1991) *Walker's Mammals of the World*. Vol.1 5th ed, The Johns Hopkins Univ Press, Baltimore and London
- Oshida T and Obara Y (1991) Karyotypes and chromosome banding patterns of a male Japanese giant flying squirrel, *Petaurista leucogenys* TEMMINCK. *Chrom Inf Serv* 50: 26–28
- Oshida T and Obara Y (1993) C-band variation in the chromosomes of the Japanese giant flying squirrel, *Petaurista leucogenys*. *J Mammal Soc Jpn* 18: 61–67
- Oshida T and Yoshida MC (1999a) Chromosomal characterization

- and karyotypic evolution of some Asian squirrels. *Jpn J Zoo Wildl Med* 4: 135–142 (in Japanese with English abstract)
- Oshida T and Yoshida MC (1999b) Chromosomal localization of nucleolus organizer regions in eight Asian squirrel species. *Chrom Sci* 3: 55–58
- Saccone C, Pesole G, Sbisà E (1991) The main regulatory region of mammalian mitochondrial DNA: structure-function model and evolutionary pattern. *J Mol Evol* 33: 83–91
- Saitou N, Nei M (1987) The neighbor-joining method: A new method reconstructing phylogenetic trees. *Mol Biol Evol* 4: 406–425
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular Cloning: A Laboratory Manual*. 2nd ed, Cold Spring Harbor Laboratory, New York
- Shikama T (1949) The Kuzuü, Ossuaries. Geological and palaeontological studies of the limestone fissure deposits, in Kuzuü, Totigi Prefecture. *Sci Rep Tohoku Univ 2nd Ser* 23: 1–209
- Southern SO, Southern PJ, Dizon AE (1988) Molecular characterization of a cloned dolphin mitochondrial genome. *J Mol Evol* 28: 32–42
- Swofford DL (1993) *User Manual for PAUP Version 3.1: Phylogenetic analysis using parsimony*. Illinois Natural History Survey, Champaign, Illinois
- Thompson JD, Higgins DG, T. Gibson J (1994) Clustal W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* 22: 4673–4680
- Vigilant L, Pennington R, Harpending H, Kocher TD, Wilson AC (1989) Mitochondrial DNA sequences in single hairs from a southern African population. *Proc Natl Acad Sci USA* 86: 9350–9354

(Received June 8, 2000 / Accepted September 11, 2000)