



Serological and molecular surveys of babesia bovis and babesia bigemina among native cattle and cattle imported from thailand in hue, Vietnam

著者 (英)	Sivakumar Thillaiampalam, Lan Dinh Thi Bich, Long Phung Thang, Viet Le Quoc, Weerasooriya Gayani, Kume Aiko, Suganuma Keisuke, Igarashi Ikuo, Yokoyama Naoaki
journal or publication title	Journal of Veterinary Medical Science
volume	80
number	2
page range	333-336
year	2018
URL	http://id.nii.ac.jp/1588/00004250/

doi: info:doi/10.1292/jvms.17-0549



NOTE

Parasitology

Serological and molecular surveys of *Babesia bovis* and *Babesia bigemina* among native cattle and cattle imported from Thailand in Hue, Vietnam

Thillaiampalam SIVAKUMAR¹⁾, Dinh Thi Bich LAN²⁾, Phung Thang LONG³⁾,
Le Quoc VIET²⁾, Gayani WEERASOORIYA^{1,4)}, Aiko KUME¹⁾,
Keisuke SUGANUMA^{1,5)}, Ikuo IGARASHI¹⁾ and Naoaki YOKOYAMA^{1)*}

¹⁾National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary Medicine, Inada-cho, Obihiro, Hokkaido 080-8555, Japan

²⁾Hue University Institute of Biotechnology, Phu Thuong Commune, Phu Vang District, Thua Thien Hue Province 47000, Vietnam

³⁾University of Agriculture and Forestry, Hue University, 102 Phung Hung Street, Hue 47000, Vietnam

⁴⁾Veterinary Research Institute, P.O. Box 28, Peradeniya, Sri Lanka

⁵⁾Research Center for Global Agromedicine, Obihiro University of Agriculture and Veterinary Medicine, Inada, Obihiro, Hokkaido 080-8555, Japan

ABSTRACT. Serum and DNA from blood samples collected from Vietnamese yellow cattle (n=101) and cattle imported from Thailand (n=54) at a Vietnamese slaughter house were screened for *Babesia bovis* and *Babesia bigemina* infections by enzyme-linked immunosorbent assay (ELISA) and PCR. The positive rates determined by ELISA (*B. bovis* and *B. bigemina*) or PCR (*B. bigemina*) in the Vietnamese cattle were significantly higher than those found in Thai cattle. Some PCR-positive Vietnamese animals were ELISA-negative, whereas all PCR-positive Thai cattle were ELISA-positive, suggesting that the animals were infected in Thailand. Importing *Babesia*-infected cattle may lead to the introduction of new parasite strains, possibly compromising the development of anti-*Babesia* immune control strategies in Vietnam.

KEY WORDS: *Babesia bigemina*, *Babesia bovis*, cattle, Thailand, Vietnam

J. Vet. Med. Sci.

80(2): 333–336, 2018

doi: 10.1292/jvms.17-0549

Received: 10 October 2017

Accepted: 26 November 2017

Published online in J-STAGE:

15 December 2017

Hemoprotozoan parasites that infect cattle generally include species of *Babesia*, *Theileria* and *Trypanosoma*. Among bovine *Babesia* parasites, *Babesia bovis* and *Babesia bigemina*, which are economically significant pathogens, are widespread in tropical and sub-tropical regions worldwide [3]. Both parasite species are transmitted by tick vectors and cause severe clinical diseases in susceptible cattle [16]. *Babesia* sporozoites injected by the infected tick vector during its blood meal invade bovine red blood cells (RBCs) and then transform into merozoites that multiply asexually [12]. Merozoite egress from infected RBCs causes massive destruction of these cells within the blood vessels, leading to anemia and anemia-related clinical signs [3]. In addition, acute infection with *B. bovis* may result in respiratory and neurological syndromes via sequestration of infected RBCs in the capillary beds [9]. In both cases, the prognosis is poor when treatment of the disease is delayed.

Vietnam is an agricultural country. Although the livestock sector has the potential to greatly contribute to the Vietnamese national economy, the productivity of this industry remains low for various reasons, such as outbreaks of infectious diseases. Recent studies reported the detection of different species of hemoprotozoan parasites, including *B. bovis* and *B. bigemina*, in Vietnamese cattle and buffalo bred in different geographical regions of the country [17, 22, 24]. As the meat production is not sufficient to meet the domestic demand, Vietnam has recently started to import live cattle from several countries, including Thailand, for use in the meat industry. Notably, various hemoprotozoan parasite species, including *B. bovis* and *B. bigemina*, have been reported in cattle bred in Thailand [1, 7, 11, 13, 19]. However, the animals imported from Thailand have not been investigated for infections with *Babesia* parasites in Vietnam. Therefore, the objective of the present study was to investigate whether *B. bovis* and *B. bigemina* are present among cattle imported from Thailand by enzyme-linked immunosorbent assay (ELISA) systems and PCR assays.

Previously, ELISA systems based on recombinant forms of the rhoptry-associated protein-1 (RAP-1) were found to be useful

*Correspondence to: Yokoyama, N.: yokoyama@obihiro.ac.jp

©2018 The Japanese Society of Veterinary Science



This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License. (CC-BY-NC-ND 4.0: <https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Table 1. PCR primers used in the present study

Purpose	Parasite	Target gene	Forward	Reverse	Amplicon size (bp)	Reference
Sequencing	<i>B. bovis</i>	RAP-1	AAGTTTCATCGAGGATACTAACC	TCAGAGGTATCCGGGGTGTGTC	741	This study
	<i>B. bigemina</i>	RAP-1	TTGATTATGAAAGCACGTCCTC	TTACCGCATCTGAATCATCTG	522	This study
Protein expression ^{a)}	<i>B. bovis</i>	RAP-1	gggatccAACTATCTGAAAGCCAATG	ggcctegagtcAAGCAATATCTCGCCTAGG	300	[20]
	<i>B. bigemina</i>	RAP-1	gggatccCCTCACTACCTTTCTAAGGC	ggcctegagtcAATCTTCAATTTTGGGGTCAATC	300	[20]
Diagnostic PCR	<i>B. bovis</i>	RAP-1	CACGAGCAAGGAACCTACCGATGTTGA	CCAAAGGACCTTCAACGTACGAGGTCA	356	[10]
	<i>B. bigemina</i>	AMA-1	TACTGTGACGAGGACGGGATC	CCTCAAAAAGCAGATTCGGAGT	211	[21]

a) Uppercase letters in the primer sequences indicate the regions corresponding to the template sequences. The restriction sites in the forward (*Bam*HI) and reverse (*Xho*I) primers are underlined.

Table 2. ELISA and PCR results summary

Animal type	No. of samples	<i>B. bovis</i>		<i>B. bigemina</i>	
		ELISA	PCR	ELISA	PCR
Vietnamese yellow cattle	101	No. positive 74	No. positive 16	No. positive 78	No. positive 31
		% (CI) 73.3 (63.9–80.9)	% (CI) 15.8 (10–24.2)	% (CI) 77.2 (68.1–84.3)	% (CI) 30.7 (22.5–40.3)
Thai cattle	54	No. positive 23	No. positive 3	No. positive 30	No. positive 4
		% (CI) 42.6 (30.3–55.9)	% (CI) 5.6 (1.9–15.1)	% (CI) 55.6 (42.4–68.0)	% (CI) 7.4 (2.9–17.6)

CI, confidence interval.

for serological detection of *B. bovis* and *B. bigemina* infections [4, 5]. In the present study, genetic variations present in the *B. bovis* and *B. bigemina rap-1* sequences from Vietnam were analyzed before the development of RAP-1 antigen-based ELISAs. *Babesia bovis*- and *B. bigemina*-positive archived blood DNA samples sourced from Vietnamese cattle (*B. bovis*, n=8; *B. bigemina*, n=18) and buffalo (*B. bovis*, n=3; *B. bigemina*, n=2) [27] were used to amplify 741-base pair (bp) and 522-bp fragments of the N-terminal region of *rap-1* of *B. bovis* and *B. bigemina*, respectively, using the primer sets shown in Table 1. The gene fragments were then cloned into a PCR 2.1 vector (TOPO, Invitrogen, Carlsbad, CA, U.S.A.) and sequenced on an ABI PRISM 3700 genetic analyzer. The newly determined *B. bovis* and *B. bigemina rap-1* sequences were registered in GenBank (*B. bovis*, LC323157–LC323167; *B. bigemina*, LC323168–LC323188). Sequencing analyses revealed that the identity and similarity scores shared among the *rap-1* sequences of *B. bovis* (98.2–100 and 96.3–100%, respectively) and *B. bigemina* (97.5–100 and 96.5–100%, respectively) were very high, indicating that Vietnamese *B. bovis* and *B. bigemina rap-1* are highly conserved.

Subsequently, 300-bp gene fragments encoding 100 amino acids within the *B. bovis*- and *B. bigemina*-specific regions of *rap-1* sequences [4, 5] were PCR-amplified using the previously described primer sets and the PCR 2.1 vectors containing inserts from the Vietnamese *B. bovis* (LC323159) and *B. bigemina rap-1* (LC323168) gene sequences (Table 1) [20]. The PCR products were ligated into a pGEX-4T1 (*B. bovis*) or pGEX-6p2 (*B. bigemina*) plasmid vector (GE Healthcare, Little Chalfont, U.K.), and recombinant RAP-1 antigens were expressed as glutathione *S*-transferase-fusion proteins as described previously [20]. Finally, the glutathione *S*-transferase tag was cleaved using thrombin and PreScission protease (GE Healthcare) to purify the *B. bovis* and *B. bigemina* recombinant RAP-1 proteins (rRAP-1), respectively [20]. These rRAP-1 antigens were then used to develop ELISAs for the sero-diagnosis of *B. bovis* and *B. bigemina* infections. Briefly, ELISA plates were coated with 100 μ l of 1 μ g/ml *B. bovis* or *B. bigemina* rRAP-1 antigen in carbonate-bicarbonate buffer (Sigma-Aldrich, St. Louis, MO, U.S.A.) at 4°C overnight. The plates were then washed once with phosphate-buffered saline containing 0.5% Tween 20 (PBST), and then blocked with 1% bovine serum albumin (BSA; Sigma-Aldrich) (100 μ l per well) at 37°C for 1 hr. After washing once with PBST, 100 μ l of each serum sample diluted 1:100 in 1% BSA was added to the wells in duplicate, and the plate was incubated at 37°C for 1 hr. The ELISA plates were washed six times with PBST, and 100 μ l of horseradish peroxidase-conjugated rabbit anti-bovine IgG (Bethyl Laboratories, Inc., Montgomery, TX, U.S.A.) diluted 1:4,000 in 1% BSA was added to each well. The ELISA plates were incubated at 37°C for 1 hr and washed six times with PBST. Next, 50 μ l of 3,3',5,5'-tetramethylbenzidine (TMB) substrate (Sigma-Aldrich) was added to each well. The plates were incubated in the dark for 10 min, and the reaction was stopped by adding 50- μ l of TMB stop solution (Sigma-Aldrich). Finally, the optical density (OD) value was measured at 450 nm. A sample was considered positive if the OD value was higher than the cut-off value that had been determined as the sum of the mean OD values and 5 \times standard deviations of the five negative serum samples used in each plate (data not

shown) [20]. Basic Local Alignment Search Tool (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) analysis showed that the Vietnamese *B. bovis* and *B. bigemina rap-1* sequences shared 98 and 97% minimum identity scores with the sequences reported globally. Therefore, although the ELISAs in the present study were developed based on gene sequences isolated in Vietnam, they can also detect specific antibodies against *B. bovis* and *B. bigemina* of Thai origin.

Blood samples were collected from September to December (2016) at a slaughter house located in Hue from 101 Vietnamese yellow cattle, including 93 males and 8 females, and 54 Brahman cattle (males) that had been imported from Thailand. All animals were apparently healthy during sampling. The Vietnamese animals were 1–6 years old (98 and 3 animals were 1–3.5 and 4–6 years old, respectively), while the Thai cattle were 1.5–3.5 years old. The period between the departure of cattle from Thailand and time of sampling was one week. Blood samples were subjected to serum separation and DNA extraction. Serum samples were analyzed by *B. bovis*- and *B. bigemina*-specific ELISA, while DNA samples were subjected to *B. bovis*- and *B. bigemina*-specific PCR assays [10, 21] as described previously (Table 1) [26]. The positive rates were analyzed by OpenEpi software (<http://www.openepi.com/Proportion/Proportion.htm>) to determine the 95% confidence intervals using a Wilson score interval [25]. The *P* values were calculated using an “N-1” χ^2 test [6, 18] (https://www.medcalc.org/calccomparison_of_proportions.php). The rate difference was considered statistically significant when the *P* value was <0.05.

The ELISA positivity rates for *B. bovis* and *B. bigemina* in Vietnamese cattle (73.3 and 77.2%, respectively) were significantly higher than those in Thai cattle (42.6 and 55.6%, respectively) (Table 2). The *B. bigemina* PCR-positivity rate was also higher for Vietnamese cattle (30.7%) than for Thai cattle (7.4%), but a significant difference was not found for *B. bovis* PCR-positive rates between Vietnamese and Thai cattle (15.8 and 5.6%, respectively) (Table 2). Co-infections with *B. bovis* and *B. bigemina* were detected by ELISA and PCR in the parasite-infected Vietnamese (61 and 7 animals, respectively) and Thai cattle (17 and 1 animals, respectively). Of the 16 samples that were PCR-positive for *B. bovis* in Vietnamese cattle, 12 were positive by ELISA, whereas four samples that were positive by PCR were negative by ELISA. Similarly, of the 31 samples that were PCR-positive for *B. bigemina*, 22 were positive by ELISA, whereas nine PCR-positive samples were ELISA-negative. The PCR-positive but ELISA-negative results for these samples indicate that the animals likely became infected very recently and that the samples were collected before ELISA-detectable antibodies had been developed. In contrast, all Thai cattle that were *B. bovis*- and *B. bigemina*-positive by PCR were also positive by ELISA, suggesting that no recent infections occurred among Thai cattle. A previous study found that *B. bovis* antibodies in experimentally infected cattle were detectable by ELISA from days 10 to 14 post-infection [8]. In the present investigation, Thai animals were sampled one week after they left Thailand. Thus, the Thai cattle were likely infected before importation. However, additional studies are needed to test the animals for *Babesia* infections in Thailand and Vietnamese quarantine facilities before and after their importation to confirm this hypothesis.

Although the infection rates for *B. bovis* and *B. bigemina* were higher among Vietnamese cattle compared to in cattle imported from Thailand, the introduction of new parasite strains may complicate the development of immune control strategies in Vietnam. For example, genetic analyses of *B. bovis* merozoite surface antigen-1 (*msa-1*) demonstrated that the genotypic distribution of *B. bovis* differs between Vietnam and Thailand [15, 23, 27]. Previous studies showed that the immune profiles of *B. bovis*-infected cattle may differ in a strain-specific manner based on the genetic variation observed in merozoite surface antigens, such as *msa-1* [2, 14]. Thai cattle typically arrive in Hue 1–3 weeks after departing from Thailand. The animals graze occasionally during transportation and before slaughtering. Therefore, tick vectors in Vietnamese pasture may acquire *Babesia* infection from Thai cattle during blood meal. In addition, entry of *Babesia*-infected ticks together with imported cattle into Vietnam cannot be ruled out.

In conclusion, the present study, which analyzed *B. bovis* and *B. bigemina* infections in Vietnam using molecular and serological diagnostic tools, found that cattle imported from Thailand had been exposed to and still harbored these parasite species. Therefore, Thai cattle should be tested for *B. bovis* and *B. bigemina* positivity before importing them into Vietnam to ensure that only the *Babesia*-free animals are imported into the country.

ACKNOWLEDGMENTS. We thank Ms. Hiroko Yamamoto, National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary Medicine, for her excellent technical assistance. This study was supported by grants from the Japan Society for the Promotion of Science (JSPS) KAKENHI (grant numbers: 16H05033, 15K14862 and 26257417), Open Partnership Joint Projects of the JSPS Bilateral Joint Research Projects, and AMED/JICA Science and Technology Research Partnership for Sustainable Development (SATREPS).

REFERENCES

1. Altangerel, K., Sivakumar, T., Inpankaew, T., Jittapalpong, S., Terkawi, M. A., Ueno, A., Xuan, X., Igarashi, I. and Yokoyama, N. 2011. Molecular prevalence of different genotypes of *Theileria orientalis* detected from cattle and water buffaloes in Thailand. *J. Parasitol.* **97**: 1075–1079. [Medline] [CrossRef]
2. Berens, S. J., Brayton, K. A., Molloy, J. B., Bock, R. E., Lew, A. E. and McElwain, T. F. 2005. Merozoite surface antigen 2 proteins of *Babesia bovis* vaccine breakthrough isolates contain a unique hypervariable region composed of degenerate repeats. *Infect. Immun.* **73**: 7180–7189. [Medline] [CrossRef]
3. Bock, R., Jackson, L., de Vos, A. and Jorgensen, W. 2004. Babesiosis of cattle. *Parasitology* **129** Suppl: S247–S269. [Medline] [CrossRef]
4. Boonchit, S., Xuan, X., Yokoyama, N., Goff, W. L., Waghela, S. D., Wagner, G. and Igarashi, I. 2004. Improved enzyme-linked immunosorbent assay using C-terminal truncated recombinant antigens of *Babesia bovis* rhoptry-associated protein-1 for detection of specific antibodies. *J. Clin. Microbiol.* **42**: 1601–1604. [Medline] [CrossRef]

5. Boonchit, S., Alhassan, A., Chan, B., Xuan, X., Yokoyama, N., Ooshiro, M., Goff, W. L., Waghela, S. D., Wagner, G. and Igarashi, I. 2006. Expression of C-terminal truncated and full-length *Babesia bigemina* rhoptyry-associated protein 1 and their potential use in enzyme-linked immunosorbent assay. *Vet. Parasitol.* **137**: 28–35. [[Medline](#)] [[CrossRef](#)]
6. Campbell, I. 2007. Chi-squared and Fisher-Irwin tests of two-by-two tables with small sample recommendations. *Stat. Med.* **26**: 3661–3675. [[Medline](#)] [[CrossRef](#)]
7. Cao, S., Aboge, G. O., Terkawi, M. A., Yu, L., Kamyngkird, K., Luo, Y., Li, Y., Goo, Y. K., Yamagishi, J., Nishikawa, Y., Yokoyama, N., Suzuki, H., Igarashi, I., Maeda, R., Inpankaew, T., Jittapalpong, S. and Xuan, X. 2012. Molecular detection and identification of *Babesia bovis* and *Babesia bigemina* in cattle in northern Thailand. *Parasitol. Res.* **111**: 1259–1266. [[Medline](#)] [[CrossRef](#)]
8. Chung, C. J., Suarez, C. E., Bandaranayaka-Mudiyanselage, C. L., Bandaranayaka-Mudiyanselage, C. B., Rzepka, J., Heiniger, T. J., Chung, G., Lee, S. S., Adams, E., Yun, G. and Waldron, S. J. 2017. A novel modified-indirect ELISA based on spherical body protein 4 for detecting antibody during acute and long-term infections with diverse *Babesia bovis* strains. *Parasit. Vectors* **10**: 77. [[Medline](#)] [[CrossRef](#)]
9. Everitt, J. I., Shaddock, J. A., Steinkamp, C. and Clabaugh, G. 1986. Experimental *Babesia bovis* infection in Holstein calves. *Vet. Pathol.* **23**: 556–562. [[Medline](#)] [[CrossRef](#)]
10. Figueroa, J. V., Chieves, L. P., Johnson, G. S. and Buening, G. M. 1993. Multiplex polymerase chain reaction based assay for the detection of *Babesia bigemina*, *Babesia bovis* and *Anaplasma marginale* DNA in bovine blood. *Vet. Parasitol.* **50**: 69–81. [[Medline](#)] [[CrossRef](#)]
11. Garcia, H. A., Kamyngkird, K., Rodrigues, A. C., Jittapalpong, S., Teixeira, M. M. and Desquesnes, M. 2011. High genetic diversity in field isolates of *Trypanosoma theileri* assessed by analysis of cathepsin L-like sequences disclosed multiple and new genotypes infecting cattle in Thailand. *Vet. Parasitol.* **180**: 363–367. [[Medline](#)] [[CrossRef](#)]
12. Homer, M. J., Aguilar-Delfin, I., Telford, S. R. 3rd., Krause, P. J. and Persing, D. H. 2000. Babesiosis. *Clin. Microbiol. Rev.* **13**: 451–469. [[Medline](#)] [[CrossRef](#)]
13. Kashiwazaki, Y., Pholpark, M., Polsar, C. and Pholpark, S. 1998. Haemoparasite infections in newly introduced dairy cattle in Loei Province, Thailand: *Trypanosoma evansi* antigen levels by ELISA referring to abortion. *Vet. Parasitol.* **80**: 99–109. [[Medline](#)] [[CrossRef](#)]
14. Leroith, T., Brayton, K. A., Molloy, J. B., Bock, R. E., Hines, S. A., Lew, A. E. and McElwain, T. F. 2005. Sequence variation and immunologic cross-reactivity among *Babesia bovis* merozoite surface antigen 1 proteins from vaccine strains and vaccine breakthrough isolates. *Infect. Immun.* **73**: 5388–5394. [[Medline](#)] [[CrossRef](#)]
15. Liyanagunawardena, N., Sivakumar, T., Kothalawala, H., Silva, S. S., Battsetseg, B., Lan, D. T. B., Inoue, N., Igarashi, I. and Yokoyama, N. 2016. Type-specific PCR assays for *Babesia bovis* *msa-1* genotypes in Asia: Revisiting the genetic diversity in Sri Lanka, Mongolia, and Vietnam. *Infect. Genet. Evol.* **37**: 64–69. [[Medline](#)] [[CrossRef](#)]
16. Mosqueda, J., Olvera-Ramirez, A., Aguilar-Tipacamú, G. and Canto, G. J. 2012. Current advances in detection and treatment of babesiosis. *Curr. Med. Chem.* **19**: 1504–1518. [[Medline](#)] [[CrossRef](#)]
17. Nguyen, T. T., Zhou, M., Ruttayaporn, N., Nguyen, Q. D., Nguyen, V. K., Goto, Y., Suzuki, Y., Kawazu, S. and Inoue, N. 2014. Diagnostic value of the recombinant tandem repeat antigen TeGM6-4r for surra in water buffaloes. *Vet. Parasitol.* **201**: 18–23. [[Medline](#)] [[CrossRef](#)]
18. Richardson, J. T. 2011. The analysis of 2×2 contingency tables—yet again. *Stat. Med.* **30**: 890, author reply 891–892. [[Medline](#)] [[CrossRef](#)]
19. Simking, P., Saengow, S., Bangphoomi, K., Sarataphan, N., Wongnarkpet, S., Inpankaew, T., Jittapalpong, S., Munkhjargal, T., Sivakumar, T., Yokoyama, N. and Igarashi, I. 2013. The molecular prevalence and MSA-2b gene-based genetic diversity of *Babesia bovis* in dairy cattle in Thailand. *Vet. Parasitol.* **197**: 642–648. [[Medline](#)] [[CrossRef](#)]
20. Sivakumar, T., Kothalawala, H., Weerasooriya, G., Silva, S. S. P., Puvanendiran, S., Munkhjargal, T., Igarashi, I. and Yokoyama, N. 2016. A longitudinal study of *Babesia* and *Theileria* infections in cattle in Sri Lanka. *Vet. Parasitol. Reg. Stud. Rep* **6**: 20–27.
21. Sivakumar, T., Altangerel, K., Battsetseg, B., Battur, B., Aboulaila, M., Munkhjargal, T., Yoshinari, T., Yokoyama, N. and Igarashi, I. 2012. Genetic detection of *Babesia bigemina* from Mongolian cattle using apical membrane antigen-1 gene-based PCR assay. *Vet. Parasitol.* **187**: 17–22. [[Medline](#)] [[CrossRef](#)]
22. Sivakumar, T., Lan, D. T. B., Long, P. T., Yoshinari, T., Tattiyapong, M., Guswanto, A., Okubo, K., Igarashi, I., Inoue, N., Xuan, X. and Yokoyama, N. 2013. PCR detection and genetic diversity of bovine hemoprotozoan parasites in Vietnam. *J. Vet. Med. Sci.* **75**: 1455–1462. [[Medline](#)] [[CrossRef](#)]
23. Tattiyapong, M., Sivakumar, T., Takemae, H., Simking, P., Jittapalpong, S., Igarashi, I. and Yokoyama, N. 2016. Genetic diversity and antigenicity variation of *Babesia bovis* merozoite surface antigen-1 (MSA-1) in Thailand. *Infect. Genet. Evol.* **41**: 255–261. [[Medline](#)] [[CrossRef](#)]
24. Weerasooriya, G., Sivakumar, T., Lan, D. T. B., Long, P. T., Takemae, H., Igarashi, I., Inoue, N. and Yokoyama, N. 2016. Epidemiology of bovine hemoprotozoa parasites in cattle and water buffalo in Vietnam. *J. Vet. Med. Sci.* **78**: 1361–1367. [[Medline](#)] [[CrossRef](#)]
25. Wilson, E. B. 1927. Probable inference, the law of succession, and statistical inference. *J. Am. Stat. Assoc.* **22**: 209–212. [[CrossRef](#)]
26. Ybañez, A. P., Sivakumar, T., Ybañez, R. H., Vincoy, M. R., Tingson, J. A., Perez, Z. O., Gabotero, S. R., Buchorno, L. P., Inoue, N., Matsumoto, K., Inokuma, H. and Yokoyama, N. 2013. Molecular survey of bovine vector-borne pathogens in Cebu, Philippines. *Vet. Parasitol.* **196**: 13–20. [[Medline](#)] [[CrossRef](#)]
27. Yokoyama, N., Sivakumar, T., Tuvshintulga, B., Hayashida, K., Igarashi, I., Inoue, N., Long, P. T. and Lan, D. T. B. 2015. Genetic variations in merozoite surface antigen genes of *Babesia bovis* detected in Vietnamese cattle and water buffaloes. *Infect. Genet. Evol.* **30**: 288–295. [[Medline](#)] [[CrossRef](#)]