



Length of lags in responses of milk yield and somatic cell score on test day to heat stress in Holsteins

著者 (英)	Hagiya Koichi, Bamba Ikumi, Osawa Takefumi, Atagi Yamato, Takusari Naozumi, Itoh Fumiaki, Yamazaki Takeshi
journal or publication title	Animal Science Journal
volume	90
number	5
page range	613-618
year	2019-05-01
URL	http://id.nii.ac.jp/1588/00004378/

doi: info:doi/10.1111/asj.13186

1 **Length of lags in responses of milk yield and somatic cell score**
2 **on test day to heat stress in Holsteins**

3

4 Authors: Koichi HAGIYA¹, Ikumi BAMBA¹, Takefumi OSAWA², Yamato ATAGI³,
5 Naozumi TAKUSARI⁴, Fumiaki ITOH⁴ and Takeshi YAMAZAKI⁴

6

7 Institute, address, country: ¹*Obihiro University of Agriculture and Veterinary*
8 *Medicine, Department of Life and Food Science, Obihiro 080-8555,*
9 *Japan, ²National Livestock Breeding Center, Data Analysis Division, Nishigo,*
10 *Fukushima 961-8511, Japan, ³ Department of Agricultural and Environmental*
11 *Biology, Graduate School of Agricultural and Life Sciences, The University of*
12 *Tokyo, Tokyo 113-8657, Japan, ⁴NARO Hokkaido Agricultural Research Center,*
13 *Sapporo 062-8555, Japan*

14

15 Running head: Lags in heat stress response in cows

16

17 Corresponding author: Koichi Hagiya, E-mail: hagiya@obihiro.ac.jp

18

1 **Abstract**

2 We used daily records from provincial Japanese weather stations and monthly
3 test-day records of milk production to investigate the length of the lags in the
4 responses of cows' milk yield and somatic cell score (SCS) to heat stress (HS).
5 We also investigated HS thresholds in milk yield and SCS. Data were a total of
6 17 245 709 test-day records for milk and SCS in Holstein cows that had calved
7 for the first time between 2000 and 2015, along with weather records from 60
8 weather stations. Temperature–humidity index (THI) values were estimated by
9 using average daily temperature and average daily relative humidity. Adjusted
10 THI values were calculated by using temperature, relative humidity, wind speed,
11 and solar radiation. The model contained herd, calving year, month of test day,
12 age group, days in milk, and THI as a fixed effect. THIs for each day from 14
13 days before the test day until the test day were used to represent HS effects. HS
14 occurring 3 days, and between 8 and 10 days, before the test day had the
15 greatest effect on milk yield and SCS, respectively. The threshold THI values for
16 HS effect were about 60 to 65 for both traits.

17 **Keywords:** heat stress, Holstein, temperature–humidity index (THI), test day

18

1 **Introduction**

2 Negative effects of heat stress (HS) in dairy cows have been widely
3 investigated because of economic losses in the dairy industry globally. HS
4 affects feed intake, yield, and reproduction (e.g., Hayes *et al.*, 2003; West, 2003).
5 Public weather stations supply useful information for studies of HS in dairy cattle
6 (Ravagnolo *et al.*, 2000). The temperature–humidity index (THI) is generally
7 used to estimate the effects of HS. Test-day milk yield decreases by about 0.2 kg
8 per unit increase in THI (Ravagnolo *et al.*, 2000). A high THI has been
9 associated with increased somatic cell score (SCS) in several studies
10 (Hammami *et al.*, 2013; Lambertz *et al.*, 2014). Hammami *et al.* (2013) evaluated
11 indices of the effects of HS on milk, fat, protein, and SCS; they reported the
12 superiority of using a THI that was adjusted for wind speed and solar radiation,
13 as reported by Mader *et al.* (2006).

14 Heat stress in Holstein cattle is an issue of growing concern for the dairy
15 industry in Japan. Nagamine and Sasaki (2008) investigated the effects of
16 temperature on fertility in Japan. They found that temperature had highly
17 significant negative effects on conception rates in southern Japan. Hagiya *et al.*
18 (2017) reported that negative seasonal effects on SCS and conception ratio at

1 first service were larger in southern Japan than in central and northern Japan.
2 Atagi *et al.* (2017) reported seasonal changes in semen production traits with
3 changes in temperature and humidity data from national weather stations.

4 In previous studies, a delayed response of test-day yield to HS has been
5 reported. Hammami *et al.* (2015) assigned a THI averaged over the 3 days
6 before the test day for yield traits and SCS in their analysis of HS effects in
7 Luxembourg and Germany. Bernabucci *et al.* (2014) reported that the greatest
8 negative effect on yield traits (milk, fat, and protein yields and fat and protein
9 percentages) in Italian Holsteins was observed when HS occurred 4 days before
10 the test day. Carabaño *et al.* (2016) used the average THI of the test day and the
11 2 preceding days in their analysis of HS, to take into account the length of the lag
12 in the response to HS in Holsteins in Belgium, Luxembourg, Slovenia, and
13 Spain; the results varied across countries.

14 The number of studies of SCS response to HS is limited compared with
15 those of milk yield and components. Moreover, to our knowledge, no published
16 reports have used THI to investigate the length of the lag in the response of SCS
17 to HS in Japan or northeast Asia.

18 Here, our main aim was to elucidate the lengths of the lags in responses

1 of milk yield and SCS to HS in cows by using test-day records and daily weather
2 records from provincial weather stations in Japan. We also investigated
3 threshold values of THI in milk yield and SCS.

4

5 **Materials and methods**

6 *Data*

7 Test-day records of milk and somatic cell count (SCC) at 6 through 305
8 days in milk in Holstein cows that had calved for the first time between 2000 and
9 2015 were provided by the Livestock Improvement Association of Japan (Tokyo,
10 Japan). Records were collected through the Dairy Herd Improvement Program.
11 The data included records of test-day milk yield and SCC for first-lactation cows
12 from all over Japan. The SCCs were log-transformed into SCSs by using the
13 following formula (Ali & Shook, 1980):

$$14 \quad \text{SCS} = \log_2(\text{SCC}/100\,000) + 3.$$

15 Weather records from 60 provincial weather stations for the period from
16 2000 to 2015 were obtained from the website of MeteoCrop DB (Institute for
17 Agro–Environmental Science, NARO, 2017). Ravagnolo *et al.* (2000) reported
18 that maximum daily temperature and minimum daily relative humidity were the

1 most critical variables in calculating THI to quantify HS. However, in our
2 preliminary study using weather records in Japan, we found that THI based on
3 average temperature and average relative humidity in a day was more effective
4 than that calculated by using maximum temperature and lowest relative humidity
5 in a day. Therefore, THI values were estimated by using average daily
6 temperature and average daily relative humidity. First, THI was estimated by
7 using the following formula (NRC, 1971):

$$8 \quad \text{THI} = 1.8 \times t + 32 (0.55 - 0.0055 \times rh) \times (1.8 \times t - 26), \quad [1]$$

9 where t is temperature in degrees Celsius and rh is relative humidity as a
10 percentage. In the preliminary study, we estimated the HS effect as THI adjusted
11 for a daily average wind speed and solar radiation in a day (Mader *et al.*, 2006).
12 However, the AICs with adjusted THI were similar to those with THI. That is, no
13 superiority of adjusted THI over THI was observed under Japanese weather
14 conditions.

15 A total of 17 245 709 test-day records from 2 018 406 cows were used.
16 Five subsets (divided randomly by herd because of computing memory
17 limitations) were analyzed separately in the cases of milk yield and SCS. Mean
18 daily milk yield was 27.1 kg, and mean SCS was 2.33 (Table 1). Mean THI

1 ranged from 50.7 to 51.0 (Table 2). The minimum THI was 4; the respective
 2 maximum value was 84.

3

4 *Model*

5 Test-day records were linked to the data from provincial weather stations
 6 in the 14 branch in Hokkaido, which is northern island in Japan, and in the other
 7 46 prefectures. The effects of HS were estimated by using a statistical model, as
 8 follows:

$$9 \quad y_{ijklmn} = H_i + Y_j + M_k + A_l + DIM_m + THI_n + e_{ijklmn} \quad [2]$$

10 where y_{ijklmn} is an observation of test-day milk or SCS; H_i is the fixed effect of
 11 herd i ; Y_j is the fixed effect of year at calving j (16 subclasses); M_k is the fixed
 12 effect of month k (12 calendar months); A_l is the fixed effect of age group l (18 to
 13 20, 21 and 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34 and 35 months);
 14 DIM_m is days in milk m (300 subclasses); THI_n is the index of HS as expressed
 15 by THI n (81 subclasses); and e_{ijklmn} represents vectors of random residual
 16 effects. THIs for any single day from 14 days before the test day until the test
 17 day were used to represent HS effects. When a model did not contain the fixed
 18 effects of HS, it was assumed to be a basic model. Akaike's information criterion

1 (AIC) and the least-squares (LSM) mean within each of five subsets were
 2 estimated by using the GENMOD procedure for AIC, or the GLM procedure for
 3 the LSM (SAS Institute, 2016) and compared among models with different THIs.
 4 The fitness of each subset was compared with the difference of the AIC for the
 5 basic model. Analyses were conducted separately for each of the five subsets,
 6 and means and standard errors were calculated from these results.

7 Assuming that the effect of HS was linear, the breakpoint of THI was
 8 estimated by segmented-regression analysis by using the Segmented package
 9 (Muggeo, 2003, 2008) of R (R Core Team, 2014). In a way similar to the method
 10 of Carabaño *et al.* (2016), LSMs of THI effects in equation [2] were used as the
 11 dependent variables and the slope of the segmented linear regression at lower
 12 than the breakpoint was assumed to be 0, as follows:

$$13 \quad y_i^* = c + e_i; \text{ when } x_i \leq BP, \text{ and}$$

$$14 \quad y_i^* = a + b * x_i + e_i; \text{ when } x_i > BP,$$

15 where y_i^* is the LSM of the THI effect estimated by using 3 (or 8) days before
 16 test-day in milk yield (or SCS), c is a constant, a is an intercept, b is a
 17 regression coefficient on THI x_i , and e_i is the random residual term. BP is the
 18 breakpoint, defined as the appropriate threshold value of THI when the linear

1 regression was applied to the HS effect. The number of records on THI is shown
2 in Figure 1. We used THI classes with more than 20,000 records for
3 segmented-regression analyses owing to the stability of LSM estimates (THI,
4 ranging from 18 to 82).

5

6 **Results**

7 ***Length of lags in response to HS***

8 When we used THI to model the effect of HS, the estimated AIC for milk
9 yield decreased sharply from test day to 3 days before the test day and then
10 increased toward 14 days before the test day (Figure 2). For SCS, the estimated
11 AIC with HS using THI decreased gradually from the test day to 7 to 10 days
12 before the test day and then increased a little toward 14 days before the test day
13 (Figure 3).

14 ***Least squares means for effect of heat stress***

15 When the effect of HS was assumed to be linear, the estimates of THI
16 breakpoint were 70.4 for milk yield and 68.5 for SCS. When the values of THI
17 exceeded a threshold, the LSM for the effects of HS on milk yield in the 3 days
18 before the test day decreased gradually with increasing THI (Figure 4). The

1 graphed changes in the effects of HS on milk yield were quadratic, and the
2 threshold values for THI ranged from about 60 to 65 THI for milk yield. For SCS,
3 the LSM increased with increasing THI beyond a threshold (Figure 5); the
4 threshold THI values were again from about 60 to 65. LSMs were increased
5 moderately when the THI values were less than 45.

6

7 **Discussion**

8 Daily milk yield and SCS were in agreement with those recently reported
9 (26.9 kg for daily milk yield and ranging from 2.3 to 2.5 for SCS) in Holstein cows
10 in Japan (Yamazaki *et al.*, 2016; Hagiya *et al.*, 2017).

11 Bohmanova *et al.* (2008) found in their preliminary study that the
12 weather data 3 days before the test day explained more of the variability in milk
13 yield than the data on the 2 days before the test day or on the test day itself.
14 Bernabucci *et al.* (2014) reported that the greatest negative effect was observed
15 when HS occurred 4 days before the test day. West (2003) similarly reported that
16 milk yield was affected by the THI as recorded 2 days before the test day. Hayes
17 *et al.* (2003), in contrast, found in preliminary investigations of their data that the
18 THI on the test day and 1, 2, 3, and 4 days before the test day had significant

1 effects on test-day yield. Here, we found no significant differences between the
2 estimated AIC for milk yield using the THI 3 days before the test day and those
3 from 2 to 6 days before the test day. Our finding here of a critical effect of the THI
4 3 days before the test day on milk yield was thus generally in line with these
5 previous results.

6 In their analysis of HS effects, Hammami *et al.* (2015) and Santana *et al.*
7 (2016) used the THI averaged over the 3 days before the test day to examine
8 effects on yield traits and SCS. Smith *et al.* (2013) also used the averaged THI
9 for the 3 days before the test day to estimate the effect of HS on SCS in Holstein
10 and Jersey cows. Here, we found no significant differences between the AIC
11 estimated for SCS by using the THI 10 days before the test day and those from 5
12 to 14 days before the test day. However, our results showed a longer lag time for
13 SCS than milk yield, with the greatest response to HS between 7 and 10 days
14 before the test day. Dry matter intake (DMI) is sensitive to the mean air
15 temperature 2 days earlier than the test day (West, 2003), and it causes a
16 decrease in the availability of nutrients used for milk synthesis (Polsky and von
17 Keyserlingk, 2017). Maintenance energy requirements are increased during heat
18 stress because of the high energy levels required for heat loss from the body

1 during heat stress (Atrian and Shahryar, 2012). These effects lead to a negative
2 energy balance, resulting in a decrease in milk.

3 A negative effect of HS on yield traits has been found when THI reaches
4 72 (Ravagnolo *et al.* 2000). HS thresholds for a THI over 69 to 72 have been
5 reported for various milk traits (Carabaño *et al.* 2017). Hammami *et al.* (2013)
6 reported that the THI threshold value was 62 for milk yield. By using the methods
7 of Hayes *et al.* (2009), Nguyen *et al.* (2017) estimated a genetic evaluation of
8 heat tolerance with a threshold of THI 60. Bernabucci *et al.* (2014) reported that
9 the threshold varied among studies; they concluded that this variation may have
10 occurred because of variations in the methods used to detect the THI thresholds.
11 They also suggested that threshold values were affected by the type of herd
12 cooling system, such as fans or sprinklers. Carabaño *et al.* (2017) concluded
13 that estimates of thresholds can vary with climatic regions and milk production
14 levels. In recent reports, relatively low THI thresholds, ranging from 60 to 65,
15 have been reported for HS effects on yield traits (e.g., Ammer *et al.* 2016;
16 Nguyen *et al.* 2016). Our THI thresholds seemed to be about 60 and 65 for both
17 traits—similar to the values in these recent reports. However, further studies on
18 HS thresholds are needed to elucidate quadratic changes in the effect of HS on

1 milk yield. For SCS, when the THI values were less than 45, the estimated
2 effects of THI appeared higher than when the THI values were around 50 (see
3 Figure 5); the optimal THI for Holstein cows seemed to be around 50. However,
4 the differences between the effects of low THIs and moderate THIs were quite
5 small. For SCS, Bertocchi *et al.* (2014) reported that the slope ($0.0003 \times \text{THI}$) on
6 THI was small before the breakepoint. Our results were in line with their trends,
7 and the breakpoints we obtained seemed to be somewhat higher than the THI
8 thresholds. However, the breakpoints estimated here may be applicable when
9 the genetic merit of animals for heat tolerance is estimated by using a random
10 regression model (Ravagnolo & Misztal, 2000; Santana *et al.* 2016; Atagi *et al.*
11 2018).

12 We investigated the lengths of the lags in responses of milk yield and
13 SCS to HS in cows by using test-day records and daily records from provincial
14 weather stations in Japan. HS occurring 3 days before the test day had the
15 greatest effect on milk yield. For effect on SCS, there was a longer lag in the
16 response to HS: the peak response was when HS occurred between 7 and 10
17 days before the test day. The threshold THI effect values of thresholds were
18 about 60 to 65 for both traits.

1

2 **Acknowledgments**

3 This work was supported by a grant from the Ministry of Agriculture,
4 Forestry, and Fisheries of Japan (Development of Breeding Technology for
5 Animal Life Production).

6

7 **References**

- 8 Ali AKA and Shook GE 1980. An optimum transformation for somatic cell concentration
9 in milk. *Journal of Dairy Science*, 63, 487–490.
- 10 Ammer S, Lambertz C and Gauly M 2016. Is reticular temperature a useful indicator of
11 heat stress in dairy cattle? *Journal of Dairy Science*, 99, 1–10.
- 12 Atagi Y, Onogi A, Kinukawa M, Ogino A, Kurogi K, Uchiyama K, Yasumori T, Adachi K,
13 Togashi K and Iwata H 2017. Genetic analysis of semen production traits of
14 Japanese Black and Holstein bulls: genome-wide markers-based estimation of
15 genetic parameters and environmental effect trends. *Journal of Animal Science*, 95,
16 1900–1912.
- 17 Atagi Y, Onogi A, Osawa T, Yasumori T, Adachi K, Yamaguchi S, Aihara M, Gothoh H,
18 Togashi K and Iwata H 2018. Effect of heat stress on production traits of Holstein

- 1 cattle in Japan: Parameter estimation using test-day records of first parity and
2 genome-wide markers. *Interbull Bulletin*, 53, 9-16.
- 3 Atrian P and Shahryar HA 2012. Heat stress in dairy cows (A review). *Research in*
4 *Zoology*, 2, 31-37.
- 5 Bernabucci U, Biffani S, Buggiotti L, Vitali A, Lacetera N and Nardona A 2014. The
6 effects of heat stress in Italian Holstein dairy cattle. *Journal of Dairy Science*, 97,
7 471–486.
- 8 Bertocchi L, Vitali A, Lacetera N, Nardone A, Varisco G and Bernabucci U 2014.
9 Seasonal variations in the composition of Holstein cow's milk and
10 temperature–humidity index relationship. *Animal*, 8, 667-674.
- 11 Bohmanova J, Misztal I, Tsuruta S, Norman HD and Lawlor TJ 2008. Genotype by
12 environment interaction due to heat stress. *Journal of Dairy Science*, 91, 840–846.
- 13 Carabaño MJ, Logar B, Bormann J, Minet J, Vanrobays M-L, Díaz C, Tychon B, Gengler
14 N and Hammami H 2016. Modeling heat stress under different environmental
15 conditions. *Journal of Dairy Science*, 99, 3798–3814.
- 16 Carabaño MJ, Ramon M, Diaz C, Lolina A, Perez-Guzman Serradilla JM 2017.
17 Breeding for resilience to heat stress effects in dairy ruminants. A comprehensive
18 review. *Journal of Animal Science*, 95, 1813–1826.

- 1 Hagiya K, Hayasaka K, Yamasaki T, Shirai T, Osawa T, Terawaki Y, Nagamine Y,
2 Masuda Y and Suzuki M 2017. Effects of heat stress on production, somatic cell
3 score, and conception rate in Holsteins. *Animal Science Journal*, 88, 3–10.
- 4 Hammami H, Bormann J, M'hamdi N, Montalda MM and Gengler N 2013. Evaluation of
5 heat stress effects on production traits and somatic cell score of Holsteins in
6 temperate environment. *Journal of Dairy Science*, 96, 1844–1855.
- 7 Hammami H, Vandenplas J, Vanrobays M-L, Rekik B, Bastin C and Gengler N 2015.
8 Genetic analysis of heat stress effects on yield traits, udder health, and fatty acids of
9 Walloon Holstein cows. *Journal of Dairy Science*, 98, 4956–4968.
- 10 Hayes BJ, Bowman PJ, Chamberlain AC, Verbyla K and Goddard ME 2009. Accuracy
11 of genomic breeding values in multibreed dairy cattle populations. *Genetics Selection
12 Evolution* 41:51.
- 13 Hayes BJ, Carrick M, Bowman P and Goddard ME 2003. Genotype x environment
14 interaction for milk production of daughters of Australian dairy sires from test-day
15 records. *Journal of Dairy Science*, 86, 3736–3744.
- 16 Institute for Agro-Environmental Sciences, NARO 2017. MeteoCrop DB. Retrieved on
17 12 January 2018 from http://meteocrop.dc.affrc.go.jp/amedas_data.php
- 18 Lambertz C, Sanker C and Gauly M 2014. Climatic effects on milk production traits and

- 1 somatic cell score in lactating Holstein-Friesian cows in different housing systems.
2 Journal of Dairy Science, 97, 319–329.
- 3 Mader TL, Davis MS and Brown-Brandl T. 2006. Environmental factors influencing heat
4 stress in feedlot cattle. Journal of Animal Science, 84, 712–719.
- 5 Muggeo VMR 2003. Estimating regression models with unknown break-points.
6 Statistics in medicine, 22, 3055–3071.
- 7 Muggeo VMR 2008. Segmented: An R package to fit regression models with
8 broken-line relationships. R News 8/1, 20–25. Retrieved on 12 January 2018 from
9 <http://cran.r-project.org/doc/Rnews/>.
- 10 Nagamine Y and Sasaki O 2008. Effect of environmental factors on fertility of
11 Holstein–Friesian cattle in Japan. Livestock Science, 115, 89–93.
- 12 Nguyen TTT, Bowman P, Haile-Mariam M, Nieuwhof GJ and Hayes BJ 2017. Short
13 communication: Implementation of a breeding value for heat tolerance in Australian
14 dairy cattle. Journal of Dairy Science, 100, 7362–7367.
- 15 Nguyen TTT, Bowman P, Haile-Mariam M, Pryce JE and Hayes BJ 2016. Genomic
16 selection for heat tolerance in Australian dairy cattle. Journal of Dairy Science, 99,
17 2849–2862.
- 18 NRC 1971. A guide to environmental research on animals. National Academy of

- 1 Science, Washington, DC.
- 2 Polsky L and von Keyserlingk AG 2017. Invited review: Effects of heat stress on dairy
3 cattle welfare. *Journal of Dairy Science*, 100, 8645-8657. .
- 4 R Core Team 2014. R: A language and environment for statistical computing, reference
5 index version 3.1.1. R Foundation for Statistical Computing, Vienna, Austria.
6 Retrieved on 12 January 2018 from <http://www.R-project.org/>.
- 7 Ravagnolo O and Misztal I 2000. Genetic component of heat stress in dairy cattle,
8 parameter estimation. *Journal of Dairy Science*, 83, 2126–2130.
- 9 Ravagnolo O, Misztal I and Hoogenboom G 2000. Genetic Component of Heat Stress in
10 Dairy Cattle, Development of Heat Index Function. *Journal of Dairy Science* 83,
11 2120–2125.
- 12 Santana ML Jr, Bignardi AB, Pereira RJ, Stefani G and El Faro L 2016. Genetics of heat
13 tolerance for milk yield and quality in Holsteins. *Animal*, 11, 4-14.
- 14 SAS Institute 2016. SAS® 9.4 Statements: Reference, Fifth Edition. SAS Institute Inc.,
15 Cary, NC, USA.
- 16 Smith DL, Smith T, Rude BJ and Ward SH 2013. Comparison of the effects of heat
17 stress on milk and component yields and somatic cell score in Holstein and Jersey
18 cows. *Journal of Animal Science*, 96, 3028–3033.

- 1 West JW, Mullinix BG and Bernard JK. 2003. Effects of hot, humid weather on milk
- 2 temperature, dry matter intake, and milk yield of lactating dairy cows. *Journal of Dairy*
- 3 *Science*, 86, 232–242.
- 4 Yamazaki T, Hagiya K, Takeda H, Osawa T, Yamaguchi S and Nagamine Y 2016.
- 5 Effects of stage of pregnancy on variance components, daily milk yields and 305-day
- 6 milk yield in Holstein cows, as estimated by using a test-day model. *Animal*, 10,
- 7 1263–70.
- 8

Table 1 Means, standard deviations (s.d.), and minimum and maximum values for milk yield and somatic cell score

Trait/subset	N	Mean	s.d.	Minimum	Maximum
Milk yield (kg/day)					
1	3 526 655	27.1	6.5	0.2	98.0
2	3 221 502	27.0	6.6	0.2	75.9
3	3 450 506	26.9	6.3	0.2	74.7
4	3 532 269	27.2	6.4	0.2	81.8
5	3 514 777	27.2	6.4	0.2	79.0
Somatic cell score					
1	3 526 655	2.33	1.65	-3.64	11.17
2	3 221 502	2.34	1.64	-3.64	11.33
3	3 450 506	2.33	1.64	-3.64	11.90
4	3 532 269	2.33	1.65	-3.64	12.52
5	3 514 777	2.32	1.64	-3.64	11.29

Table 2 Means, standard deviations (s.d.), and minimum and maximum values for temperature–humidity index (THI) and THI adjusted for wind speed and solar radiation (THI_{adj})

Trait/subset	N	Mean	s.d.	Minimum	Maximum
THI					
1	3 526 655	51.0	15.1	4	84
2	3 221 502	50.7	15.0	4	84
3	3 450 506	50.9	15.0	4	84
4	3 532 269	51.0	15.1	4	84
5	3 514 777	51.0	15.2	4	84

1

2

1 Figure captions

2

3 Figure 1 The number of records in the temperature–humidity index (THI)

4

5 Figure 2 Estimates of Akaike's information criterion (AIC) and 95% confidence
6 intervals (bars) for milk yield, as affected by the temperature–humidity index
7 during the days before the test day

8 Basic Model, basic model without the effects of heat stress

9

10 Figure 3 Estimates of Akaike's information criterion (AIC) and 95% confidence
11 intervals (bars) for somatic cell score, as affected by the temperature–humidity
12 index during the days before the test day

13 Basic Model, basic model without the effects of heat stress

14

15 Figure 4 Least-squares means and 95% confidence intervals (bars) for the
16 effects of heat stress (as indicated by changes in the temperature–humidity
17 index during the 3 days before the test day, THI) on milk yield, and lines
18 estimated by segmented regression analysis

19

20 Figure 5 Least-squares means and 95% confidence intervals (bars) for the
21 effects of heat stress (as indicated by changes in the temperature–humidity
22 index of the 8 days before the test day, THI) on somatic cell score, and lines
23 estimated by segmented regression analysis

24

25

1 表題： ホルスタイン種の検定日記録に対する乳量および体細胞への暑熱ス
2 トレス反応における遅延期間の長さ

3
4 著者名： 萩谷功一¹・番場郁美¹・大澤剛史²・安宅倭³・田鎖直澄⁴・伊藤文彰⁴・
5 山崎武志⁴

6
7 所属： ¹帯広畜産大学生命・食料科学研究部門，帯広市 080-8555


8 ²家畜改良センター情報分析課，福島県西白河郡西郷村 961-8511

9 ³東京大学農学生命科学研究科生産・環境生物学専攻，東京都文京区
10 113-8657

11 ⁴農研機構北海道農業研究センター，札幌市豊平区 062-8555
12

13 日本国内の検定日泌乳記録および気象観測所の毎日の記録を使用し，雌牛の乳
14 量および体細胞スコア（SCS）に対する暑熱ストレス（HS）の影響が現れるま
15 での期間および閾値を調査した．データは，2000年から2015年までに初産分娩
16 したホルスタイン種の乳量およびSCSに関する17,245,709検定日の泌乳記録，
17 および60箇所の気象観測所の気象記録である．温湿度指数（THI）は，1日の
18 平均気温および平均相対湿度を使用して推定した．数学モデルには，母数効果
19 として牛群，分娩年，検定月，月齢，搾乳日数およびTHIを含めた．暑熱スト
20 レス効果の大きさを調査するため，検定日14日前から検定日までのTHIを使用
21 した．乳量およびSCSに対し，検定日3日前および8から10日前のHSの影響
22 がそれぞれ最大であった．両形質に対するHS効果の閾値は，THI60から65の
23 範囲であった．
24
25


1 Figure 1



2

3


1 Figure 2



2

3

1 Figure 3




2

3

1


2 Figure 4



3

4

1 Figure 5



2