



Relationship between metabolic hormones and ovulation of dominant follicle during the first follicular wave post-partum in high-producing dairy cows

著者 (英)	Kawashima Chiho, Fukihara Saori, Maeda Mayumi, Kaneko Etsushi, Montoya Carlos Amaya, Matsui Motozumi, Shimizu Takashi, Matsunaga Nobuyoshi, Kida Katsuya, Miyake Yoichi, Schams Dieter, Miyamoto Akio
journal or publication title	Reproduction
volume	133
number	1
page range	155-163
year	2007-01
URL	http://id.nii.ac.jp/1588/00000493/

1 **Relationship between metabolic hormones and ovulation of dominant follicle during**
2 **the first follicular wave postpartum in high-producing dairy cows**

3
4
5
6 **Chiho Kawashima¹, Saori Fukihara¹, Mayumi Maeda¹, Etsushi Kaneko¹, Carlos Amaya Montoya²,**
7 **Motozumi Matsui², Takashi Shimizu¹, Nobuyoshi Matsunaga³, Katsuya Kida⁴, Yoh-Ichi Miyake²**
8 **Dieter Schams⁵ and Akio Miyamoto¹**

9
10
11 *Short title:* The first follicular wave postpartum in dairy cows

12
13 *Key words:* Cow, Dominant follicle, First follicular wave, Metabolic hormone, Postpartum

14
15
16
17 *Graduate School of Animal and Food Hygiene¹, Department of Clinical Veterinary Science²,*
18 *Department of Agricultural and Life Science³, Field Centre of Animal Science and Agriculture⁴,*
19 *Obihiro University of Agriculture and Veterinary Medicine, Obihiro 080-8555, Japan*
20 *TU-Munich Weihenstephan, Germany⁵.*

21
22
23
24 ¹ *Correspondence.* FAX: +81-155-49-5459; e-mail: akiomiya@obihiro.ac.jp

1 **Abstract**

2

3 Recent studies suggest that insulin-like growth factor-1 (IGF-1) is a crucial regulatory factor in follicular
4 growth during early postpartum period. The aim of the present study was to determine in detail the
5 changing profiles of metabolic and reproductive hormones in relation to ovulation of the dominant follicle
6 (DF) of the first follicular wave postpartum in high-producing dairy cows. Plasma concentrations of related
7 hormones in 22 multiparous Holstein cows were measured from - 4 to 3 wk postpartum, and the
8 development of DF was observed with colour Doppler ultrasound. Thirteen cows showed ovulation by 15.2
9 d postpartum. Anovulatory cows showed higher growth hormone and lower IGF-1 levels than in ovulatory
10 cows during the peripartum period. Each DF developed similarly, and a clear blood flow in the follicle wall
11 was observed despite ovulation or anovulation. In addition, detailed endocrine profiles were analysed in 9
12 out of 22 cows. Five cows showed an increase in plasma oestradiol-17 β (E2) increase with follicular
13 growth followed by E2 peak, LH surge and ovulation. In these cows, plasma IGF-1 concentrations
14 remained high until 10 days postpartum followed by a gradual decrease. Subsequently, the insulin level
15 increased together with the E2 peak toward ovulation. These profiles were not observed in anovulatory
16 cows. In conclusion, our data strongly support the concept that IGF-1 and insulin represent ‘metabolic
17 signals’ of the resumption of ovarian function postpartum in high-producing dairy cows. Moreover, we
18 provide the first visual evidence that both ovulatory and anovulatory dominant follicle of the first follicular
19 wave postpartum are similarly supplied with active blood flow.

20

21

22

23

24

25

26

27

28

1 **Introduction**

2

3 The negative energy balance early postpartum that is characterised by the loss of body weight and
4 mobilization of body fat stores does not affect the first follicular wave postpartum (Beam & Butler 1999).
5 In fact, the medium-sized follicles appear 5 days after calving, and reach into the large follicles at 10 days
6 postpartum in most dairy cows (Savio *et al.* 1990a, b). However, almost a half of these follicles are not able
7 to ovulate, but becomes atretic or cystic. Most ovulations are observed within 3 weeks postpartum
8 (Lamming & Bulman 1976, Lucy *et al.* 1992b, Darwash *et al.* 1997). Our recent study observed that the
9 recovery of ovarian function and subsequent reproductive performance in ovulatory cows within 3 weeks
10 postpartum was superior to that of anovulatory cows (Kawashima *et al.* 2006). It is generally accepted that
11 the cow with early resumption of the ovarian function has higher fertility (Staples *et al.*, 1990; Senatore *et al.*
12 *al.*, 1996; Darwash *et al.*, 1997). However, the factors that control the first ovulation postpartum have not
13 been fully elucidated.

14 Insulin and insulin-like growth factor 1 (IGF-1), metabolic hormones that are changed by feed
15 intake, stimulate oestradiol-17 β (E2) production in the granulosa cells (Gutierrez *et al.* 1997, Glister *et al.*
16 2001, Armstrong *et al.* 2003, Butler *et al.* 2004) and the proliferation of follicular cells (Spicer *et al.* 1993,
17 Spicer & Stewart 1996). The growth of ovulatory follicles is regulated by the metabolic hormones such as
18 IGF-1 or insulin, and is susceptible to the endocrine and metabolic status during early lactation (Spicer *et al.*
19 1990, Lucy *et al.* 1992b). The liver-derived IGF-1 is a factor that regulates the final maturation of the
20 dominant follicle during the first follicular wave postpartum (Beam & Butler 1998). The circulating IGF-1
21 in ovulatory cows at the first follicular wave postpartum was higher than that in anovulatory cows (Beam &
22 Butler 1998).

23 Most of the above studies investigated a few metabolic factors during the first follicular wave
24 postpartum. Therefore, the aim of the present study was to investigate the time-dependent relationship in
25 detail among metabolic hormones including IGF-1, growth hormone (GH) and insulin, and metabolites
26 such as glucose, non-esterified fatty acids (NEFA), total cholesterol (T-cho) and aspartate aminotransferase
27 (AST), as well as the body condition score (BCS) and ovulation during the first follicular wave postpartum

1 in a homogeneous herd at early lactation period in high-producing dairy cows. In addition, we used colour
2 Doppler ultrasonography to monitor a local blood flow during the growth of individual follicles.

3 4 5 **Materials and Methods**

6
7 Experimental procedures complied with the Guide for Care and Use of Agricultural Animals of
8 Obihiro University.

9 10 ***Animals***

11 The experiment was carried out in the Field Centre of Animal Science and Agriculture, Obihiro
12 University of Agriculture and Veterinary Medicine. We used 22 multiparous Holstein cows that calved
13 between October 2004 and April 2005. We collected samples from 4 weeks prepartum to 3 weeks
14 postpartum when the period of 0-6 days after parturition was regarded as a parturient week (0 week).
15 Additionally, 9 out of 22 cows that calved between February 2005 and April 2005 were used to observe
16 the competence of the dominant follicle in detail, we collected blood samples (4times/day) from the
17 appearance of a dominant follicle (8.5 mm in diameter) to 7 days after the first ovulation (ovulatory cows)
18 or 20 days after calving (anovulatory cows). The cows were housed in a free-stall barn throughout the
19 experimental period and offered a total mixed ration consisting of grass, corn silage and concentrate. The
20 milking was performed twice daily (6:00 and 18:00). The average 305-day milk yield was approximately
21 9,700 kg.

22 23 ***Sampling***

24 Body condition score (BCS) was assessed once a week during the experimental period by the same
25 operator using a 0 to 5 scale with 0.25 intervals, where 0 = thin and 5 = very fat (Ferguson *et al.* 1994).

26 Blood samples (serum) for biochemical analysis were obtained once a week by caudal venipuncture
27 after measurement of BCS using unheparinized, silicone-coated 9 ml tubes (Venoject®, Autosep®, Gel +
28 Clot. Act., VP-AS109K,; Terumo Corporation, Japan) and the tubes were coagulated for 30 minutes at 38 C

1 in an incubator. Blood samples (plasma) for hormonal analysis were obtained by caudal venipuncture twice
2 weekly during the same period using sterile 10 ml tubes containing 200 ul of stabilizer solution (0.3 M
3 EDTA, 1 % acetyl salicylic acid, pH 7.4). In addition, in 9 out of 22 cows, we obtained blood samples 4
4 times/day at 6 hours intervals for detection (presence or absence) of LH surge to minimized the stress of
5 blood sampling 2, time/day for detection of E2 peak and 1 time/day for profile of P4 and metabolic
6 hormones. These tubes were centrifuged at 3000 rpm for 20 minutes at 4 C and the serum or plasma
7 samples were kept at -30 C until analysis for biochemical and hormonal analyses.

8 Ovaries of all cows were scanned twice weekly from 7 days to 30 days after calving by transrectal
9 ultrasonography using a colour Doppler ultrasound scanner (SSD-5500, Aloka Co., Tokyo, Japan) equipped
10 with a 7.5-MHz convex transducer (UST-995-7.5, Aloka Co.). Follicles ≥ 5 mm were measured for
11 maximum diameter and follicles ≥ 8.5 mm were regarded as dominant follicles (Ginther *et al.* 2003). After
12 morphological evaluation, the flow mode was activated for blood flow mapping and the presence or
13 absence of blood flow was assessed for each follicle. Colour signals were used to generate images in which
14 blood flow with a velocity higher than 2 mm/sec could be located as areas of colour within the follicle wall.

15 16 ***Identification of ovulation during the first follicular wave postpartum***

17 When the P4 concentration in plasma was increased to more than 1 ng/ml, the cow was confirmed as
18 having returned to luteal activity (Stevenson & Britt 1979). Furthermore, the first ovulation and luteal
19 formation were identified using an ultrasound scanner equipped with a 7.5-MHz convex transducer.

20 21 ***Measurement of P4 and E2***

22 The P4 determination in plasma was performed by enzymeimmunoassay (EIA) after extraction by
23 diethyl ether as described previously (Miyamoto *et al.* 1992), and the extraction efficiency was 93%. The
24 standard curve ranged from 0.05 to 50 ng/ml, and the ED50 of assay was 3.2 ng/ml. The intra- and
25 interassay coefficients of variation (CVs) averaged 6.7 % and 7.2 %, respectively. The E2 determination in
26 plasma was performed by EIA after extraction by diethyl ether as described previously (Acosta *et al.* 2000),
27 and the extraction efficiency was 85%. The standard curve ranged from 1.95 to 2000 pg/ml, and the ED50
28 of assay was 5.2 pg/ml. The intra- and interassay CVs averaged 6.7 % and 8.7 %, respectively.

1

2 ***Measurement of FSH, LH, GH, IGF-1 and insulin***

3 The determination for plasma FSH, LH, GH, IGF-1 and insulin was performed by EIA using the
4 biotin-streptavidin amplification technique.

5 The FSH assay was performed as previously described (Watanabe *et al.* 1997). The standard curve
6 for FSH ranged from 0.18 to 12 ng/ml, and the ED50 of the assay was 1.7 ng/ml. The intra- and interassay
7 CVs averaged 8.3 % and 14.6 %, respectively.

8 The LH assay was previously described in detail (Mutayoba *et al.* 1990). The standard curve for LH
9 ranged from 0.09 to 50 ng/ml, and the ED50 of the assay was 3.1 ng/ml. The intra- and interassay CVs
10 averaged 8.3 % and 9.4 %, respectively.

11 The GH concentration was measured by the EIA described by Roh *et al.* (1997) with slight
12 modification. A diluted rabbit antibody to bovine GH (100µl, × 150,000, D. Schams, TU-Munich
13 Weihenstephan, Germany) was distributed in all wells of a microplate coated with anti-rabbit γ-globulin
14 antiserum, then incubated for 24 h at room temperature, and the plate was decanted. After chicken serum
15 diluted in assay buffer (100µl, 1%) was added to each well, 15µl of GH standard (0.78 to 100 ng/ml,
16 NIDDK-bGH, AFP-9984C) dissolved in assay buffer or plasma was incubated in wells for 24 h. After
17 decanting the plate, biotin-labelled GH was distributed in all wells, and then incubated for 3 h. Finally,
18 colorimetric treatments were carried out. Intra- and interassay CVs were 8.1 % and 8.5 %, respectively, and
19 the ED50 in this assay system was 6.2 ng/ml.

20 The total IGF-1 determination in plasma was performed by the EIA after protein extraction by an
21 acid-ethanol (87.5% ethanol and 12.5% 2N hydrochloric acid; Daughaday *et al.* 1980) to obtain IGF-1 free
22 from binding protein. Thirty µl of human IGF-1 standard (Roche, Indianapolis, USA, 0.39 to 50 ng/ml)
23 dissolved in assay buffer or sample was added to each well coated with anti-rabbit γ-globulin antiserum. In
24 addition, 100µl of biotin-labelled hIGF-1 (× 10,000) and rabbit anti-hIGF-1 (× 40,000, NIDDK,
25 AFP18111298) diluted in assay buffer were distributed in all wells, and then incubated for 72 h at 4 C.
26 Finally, colorimetric treatments were carried out. Intra- and interassay CVs were 5.7 % and 6.6 %,
27 respectively, and the ED50 in this assay system was 2.5 ng/ml.

1 The insulin determination was carried out by EIA. Insulin standard (Sigma, Missouri, USA) was
2 diluted with charcoal-treated serum (insulin-free) at preparation. Thirty μ l of insulin standard (39 to 5000
3 pg/ml) or plasma were added to each well coated with anti-guinea pig goat γ -globulin antiserum. In
4 addition, 100 μ l of anti-bovine guinea pig insulin (\times 150,000, D. Schams) dissolved in assay buffer were
5 distributed in all wells, and then incubated for 24 h at 4 C. After decanting the plates, 100 μ l of biotin-
6 labelled bovine insulin (\times 50,000) was distributed in all wells, and then incubated for 2 h at 4 C. Finally,
7 colorimetric treatments were carried out. Intra- and interassay CVs were 9.7 % and 14.5 %, respectively,
8 and the ED50 in this assay system was 800 pg/ml.

10 ***Biochemical analyses***

11 In each sample, the concentration of glucose, NEFA, T-cho and AST were measured using a clinical
12 chemistry automated analyzer (TBA120FR, Toshiba Medical Systems Co., Tochigi, Japan).

14 ***Statistical analyses***

15 All the data from 22 cows were analyzed by repeated ANOVA. There was a tendency for a group \times
16 time interaction on plasma concentrations of E2 (P=0.09) and plasma E2 concentrations changed over time
17 in the ovulatory cows (time effect; P<0.05). Mean concentrations of E2 were calculated for each group and
18 each sampling period and significant differences were detected by Fisher's protected least significance
19 difference test.

20 For other data, no significant interaction between group and time was observed. For all metabolites,
21 metabolic hormones and BCS in 22 cows during the peripartum period, mean concentrations for each cow
22 were calculated for the prepartum or postpartum period. For the diameter of dominant follicles and
23 metabolic hormones in 22 cows at 1, 1.5 and 2 weeks postpartum, the mean diameter and mean
24 concentrations during the sampling period were calculated and analysed repeated measures ANOVA. The
25 significant differences between ovulatory and anovulatory were analysed by Student's t test.

26 In 5 cows which ovulated when we observed in detail endocrine profiles in 9 out of 22 cows, we
27 analysed metabolic hormones at the before and after E2 peak. We divided into two phases that were before

1 E2 peak (-10 to -1 days relative to E2 peak) and after E2 peak (0 to 10 days relative to E2 peak), and the
2 significant differences between two phases were analysed by paired t test.

3 Results were expressed as mean \pm standard error of mean; differences with $P < 0.05$ were considered
4 significant.

5 6 7 **Results**

8 9 *Metabolites, BCS and metabolic hormones during the peripartum period related to the ovulation of the* 10 *dominant follicle at the first follicular wave postpartum*

11 The numbers of cows showing ovulation and anovulation of a dominant follicle during the first
12 postpartum follicular wave were 13 (59.1%) and 9 (40.9%), respectively. In ovulatory cows, the interval
13 between calving and ovulation was 15.2 ± 0.8 days, and after the first ovulation, cows had an ovarian cycle
14 followed by more than 7 days of a luteal phase. The first month milk yield was 45.5 ± 0.4 kg/day in
15 ovulatory cows and 45.1 ± 0.5 kg/day in anovulatory cows.

16 Figure 1 shows the change of metabolites, BCS and metabolic hormones during the peripartum
17 period. The serum concentration of glucose was not significantly different between the ovulatory and
18 anovulatory cows during the prepartum period. Thereafter, the serum glucose concentrations of the
19 anovulatory cows was lower than that of the ovulatory cows during the postpartum period (ovulatory,
20 59.2 ± 1.3 vs. anovulatory, 52.7 ± 1.3 ; $P < 0.001$). For BCS, there was no significant difference between
21 ovulatory and anovulatory cows during the prepartum period, however, BCS in ovulatory cows tended to
22 be higher than in anovulatory cows during the postpartum period ($P = 0.08$, Fig. 1). The serum
23 concentrations of NEFA, T-cho and AST were similar between the ovulatory and the anovulatory cows
24 during the period of study (Fig. 1).

25 The plasma concentration of insulin was not significantly different between the ovulatory and
26 anovulatory cows during the pre- and postpartum period. The plasma concentration of GH in anovulatory
27 cows was higher than that of ovulatory cows during both prepartum and postpartum periods (prepartum;
28 $P < 0.01$, postpartum; $P < 0.0001$, Fig. 1). On the other hand, the plasma concentration of IGF-1 in ovulatory

1 cows was higher than that of anovulatory cows during both prepartum and postpartum periods (prepartum;
2 $P<0.05$, postpartum; $P<0.01$, Fig. 1).

3
4 ***The function and morphology of the dominant follicle during the first follicular wave postpartum, and***
5 ***metabolic status either ovulatory or anovulatory cows***

6 The dominant follicle of the first follicular wave appeared on day 7 postpartum, and the diameter of
7 the dominant follicle was approximately 10 mm and blood flow in each follicle was observed by 14 days
8 postpartum despite ovulation or anovulation (Fig. 2). In the ovulatory cows, a corpus luteum with blood
9 flow was observed by 17 days postpartum (Fig. 2). On the other hand, the dominant follicle of anovulatory
10 cows continued to grow keeping identifiable blood flow (Fig. 2).

11 The plasma concentration of FSH did not differ between the ovulatory and anovulatory cows until
12 14 days postpartum (ovulation: $2.18\pm 0.09\text{ng/ml}$ vs. anovulation: $2.05\pm 0.09\text{ng/ml}$). The ratio of the
13 existence of dominant follicle at the first scanning was 61.5% (8/13) in the ovulatory cows and 44.4% (4/9)
14 in the anovulatory cows, and the diameter of the follicle at that time was not significantly different between
15 the ovulatory and anovulatory cows (ovulation, $9.9\pm 0.7\text{mm}$ vs. anovulation, $9.9\pm 1.0\text{mm}$).

16 The diameter of the dominant follicle had similar growth in both groups (time effect; $P<0.01$, Fig. 3)
17 and did not differ between the ovulatory and the anovulatory cows during the period of study. The plasma
18 concentrations of E2 in ovulatory cows were higher at 1.5 weeks postpartum than in 1 and 2 weeks
19 postpartum ($P<0.05$, Fig. 3), and higher than that in anovulatory cows at 1.5 weeks postpartum ($P<0.01$, Fig.
20 3). However there was no significant difference between ovulatory and anovulatory cows at 2 weeks
21 postpartum, because of decrease of preovulatory E2 concentrations. In ovulatory cows, plasma GH
22 concentrations were lower ($P<0.01$) and IGF-1 concentrations were higher ($P<0.05$) compared to
23 anovulatory cows, while the plasma insulin concentrations did not differ between the ovulatory and
24 anovulatory cows during this period (Fig. 3).

25
26 ***The metabolic-endocrine factors related to ovulation of the dominant follicle during the first follicular***
27 ***wave postpartum***

1 To observe the relationship between growth of the dominant follicle, and the endocrine and
2 metabolic factors (E2, insulin, IGF-1 and GH) in detail, 9 out of 22 cows showing ovulation or anovulation
3 of dominant follicles during the first follicular wave postpartum were studied. The number of ovulatory
4 cows was 5 (55.6%), E2 peak could be confirmed on 16.0 ± 1.1 days postpartum, the interval between E2
5 peak and LH surge was approximately 21 ± 7 hours, and interval between calving and the ovulation was
6 17.0 ± 1.5 days. On the other hand, E2 peak and LH surge could not be confirmed in anovulatory cows.

7 In ovulated cows, E2 peak could be confirmed on 16.0 ± 1.1 days postpartum, and the interval
8 between E2 peak and LH surge was approximately 21 ± 7 hours (Fig. 4). On the other hand, E2 peak and LH
9 surge could not be confirmed in anovulated cows (Fig. 4). In ovulated cows, insulin concentrations during
10 and after the E2 peak period showed a strong tendency to be higher than those before the E2 peak ($P=0.06$,
11 Table 1). The GH concentrations from before until after the E2 peak did not change. However, plasma IGF-
12 1 remained high until 7 to 10 days postpartum followed by a gradual decrease, (Fig. 4). Therefore, IGF-1
13 concentration during the after E2 peak was lower than that during the before E2 peak of E2 ($P<0.05$, Table
14 1).

15 16 17 **Discussion**

18
19 The present study focused on the changes in metabolic hormones, such as IGF-1, GH and insulin,
20 occurring during the peripartum period in dairy cows and their relation to the occurrence of ovulation. The
21 results suggest that the plasma concentration of IGF-1 relates to the early growth of ovulatory dominant
22 follicle and insulin relates to the maturation and ovulation of the ovulatory dominant follicle at the first
23 follicular wave postpartum.

24 Our results showed that higher plasma GH concentrations and lower glucose and IGF-1
25 concentrations were the clear feature in anovulatory cows during the prepartum period compared to those
26 of the ovulatory cows. Previous studies have shown that the expression of the GH receptor 1A and the IGF-
27 1 mRNA in the liver declines at parturition, and GH concentrations are elevated during the postpartum
28 period, because IGF-1 concentrations decrease to low levels, reducing the negative feedback on GH

1 (Kobayashi *et al.* 1999, Pushpakumara *et al.* 2003, Radcliff *et al.* 2003). The plasma IGF-1 levels in
2 ovulatory cows at the first follicular wave postpartum were higher than that in anovulatory cows (Beam and
3 Butler, 1998). The high GH concentrations in anovulatory cows in the present study with lower IGF-1
4 concentrations during the peripartum period may have caused GH insensitivity in a liver. These findings
5 above suggest that energy status, which reflects on GH, IGF-1 and glucose levels, and ovulation are closely
6 related to each other during the early postpartum period. However, in the present study, the metabolites
7 such as NEFA, AST and T-cho did not differ between ovulatory and anovulatory cows during the
8 peripartum period, indicating that the lipid metabolism between these two groups was similar level despite
9 the different GH levels between two groups. In anovulatory cows, body fat stores might have already been
10 reduced during the peripartum period, resulting in their lower BCS than ovulatory cows. Therefore,
11 ovulatory cows may be able to handle better the negative energy balance after parturition compared to
12 anovulatory cows. The present study could not examine the exact feed intake and nutritional status, because
13 all cows were provided with feed and water *ad libitum*. Thus, future studies are necessary to investigate the
14 effect of feed intake on metabolic hormones and on the maturation and ovulation of the first postpartum
15 dominant follicle.

16 It was observed that the rate of follicular growth did not differ between ovulatory and anovulatory
17 cows during the postpartum period. The local blood flow was detected in dominant follicle of all cows
18 despite ovulation or anovulation. Angiogenesis, the formation of new network of blood vessels, is essential
19 for follicular development and ovulation (Yamada *et al.* 1994, Jiang *et al.* 2003). In addition, a recent report
20 suggested that the angiopoietin-tie system that acts on angiogenesis in concert with vascular endothelial
21 growth factor (VEGF) controls the blood vessels to maintain active angiogenesis in developing bovine
22 follicles (Hayashi *et al.* 2003). The maintenance of follicle vasculature and appropriate blood supply to the
23 large follicles is essential for follicle dominance (Acosta *et al.* 2005). However, our results showed that an
24 insufficient blood flow supply is not the cause of anovulation. The plasma E2 concentrations during this
25 period increased only in the ovulatory cows while those in the anovulatory cows did not, which are well
26 consistent with previous reports (Beam & Butler 1997, Beam & Butler 1998). Thus, it is likely that an
27 insufficient ability of granulosa cells to secrete E2 is a determinant for anovulation rather than insufficient
28 angiogenesis.

1 The plasma concentration of FSH did not differ between the ovulatory and anovulatory cows until
2 14 days postpartum. Our data confirmed the previous studies that FSH appears to be insensitive to
3 metabolic status (Beam & Butler 1998, Lamming *et al.* 1981). On the other hand, serum concentration of
4 IGF-1 during the first follicular wave postpartum were reported to be at low levels in anovulatory dairy
5 cows (Beam & Butler 1998). In the present study, the plasma IGF-1 remained elevated until 7 to 10 days
6 postpartum followed by a gradual decrease, whereas IGF-1 in the anovulatory cows was kept at a low level
7 throughout the postpartum period. The IGF-1 stimulates steroidogenesis and the proliferation in follicular
8 cells (Spicer *et al.* 1993, Spicer & Stewart 1996), and the ratio of oestrogen to progesterone in follicular
9 fluid was correlated positively with plasma IGF-1 in lactating cows (Lucy *et al.* 1992a). Also, IGF-1 works
10 with FSH synergically in promoting follicular development and E2 synthesis (Fortune *et al.* 2004). Thus,
11 the data suggest that IGF-1 is an important factor to develop the dominant follicle to the maturation stage
12 by amplifying the endocrine signal of FSH to upregulate E2 secretion from the future dominant follicle
13 postpartum.

14 The present data suggested that plasma insulin level in the ovulatory cows increased together with
15 E2 peak. In dairy cows, the serum insulin levels increase together with E2 during the development of the
16 dominant follicle (Armstrong *et al.* 2003). Insulin has been shown to stimulate E2 production in the
17 granulosa cells (Gutierrez *et al.* 1997, Glister *et al.* 2001). In addition, E2 secreted from a follicle also
18 enhances insulin secretion from the pancreas (Morimoto *et al.* 2001). These findings clearly indicate that
19 plasma insulin and E2 from follicles positively interact with each other during the development of the
20 dominant follicle. The increase in circulating insulin level by feeding modification results in advanced first
21 ovulation postpartum in the cow (Gong *et al.* 2002) and the increase in plasma LH concentration in heifer
22 (McCann & Hansel 1986). The LH pulse frequency was reported to be greater in ovulatory cows than in
23 anovulatory cows (Lamming *et al.* 1981, Canfield & Butler 1990, Jolly *et al.* 1995). Therefore, plasma
24 insulin levels closely relates to the follicle growth and ovulation by regulating LH secretion. On the other
25 hand, LH was not controlled by insulin infusion in early lactation cows (Butler *et al.* 2004). Namely, the
26 energy status resulting in high insulin levels, but not directly insulin infusion, may induce the change of
27 pulsatile LH release, follicle growth and ovulation. Consequently, the nutritional status, which leads to the
28 increase of insulin level together with E2 peak, is one of the key features of ovulatory cows. Undoubtedly,

1 further studies are needed to characterise the factors responsible for the suboptimal GnRH and LH release
2 after parturition in dairy cows.

3 As shown in Fig 5, our results demonstrated that the changes in the concentration of two metabolic
4 hormones (IGF-1 and insulin) may regulate the development of ovulatory follicle in the first follicular wave
5 postpartum. In ovulatory cows, high IGF-1 levels were maintained during growth of dominant follicle, and
6 then insulin levels increased together with the increase in E2 secretion from dominant follicle. These
7 findings indicate that IGF -1 is an essential factor for the growth of the dominant follicle, and insulin may
8 stimulate the dominant follicle to mature and reach ovulation. Collectively, it is suggested that the
9 dominant follicle, which stimulated by high IGF-1 levels during the first follicular wave postpartum,
10 become ovulatory with the increase in E2 secretion, resulting in the occurrence of LH surge and ovulation.

11 In conclusion, the present study provides convincing evidence that IGF-1 is an important factor to
12 develop the dominant follicle during the first follicular development postpartum. After that, the increase of
13 insulin level together with E2 peak may ensure the maturation and ovulation. Our findings strongly support
14 the concept that IGF-1 and insulin represent ‘metabolic signals’ of the resumption of ovarian function
15 postpartum in high-producing dairy cows. Moreover, we provide the first visual evidence that both
16 ovulatory and anovulatory dominant follicle of the first follicular wave postpartum are similarly supplied
17 with the active blood flow, suggesting that an insufficient angiogenesis is not a determinant for anovulation.

18 19 20 **Acknowledgments**

21
22 The authors thank Dr. K. Okuda, Okayama University, Japan, for P₄ antiserum, and Dr. Parlow,
23 NIDDK, for the FSH, LH and GH standard, and IGF-1 antiserum. This study was supported by the Grant-
24 in-Aid for Scientific Research of the Japan Society for the Promotion of Science (JSPS), the 21st Century
25 COE Program (A-1) of the Ministry of Education, Culture, Sports, Science, and Technology, Japan, Secure
26 and Healthy Livestock Farming Project of the Ministry of Agriculture, Forestry and Fisheries, and the
27 Japan Livestock Technology Association. C. Kawashima and E. Kaneko are supported by the COE
28 program.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

References

Acosta TJ, Ozawa T, Kobayashi S, Hayashi K, Ohtani M, Kraetzel WD, Sato K, Schams D & Miyamoto A 2000 Perioovulatory changes in the local release of vasoactive peptides, prostaglandin F_{2α} and steroid hormones from bovine mature follicles in vivo. *Biology of Reproduction* **63** 1253-1261.

Acosta TJ, Hayashi KG, Matsui M & Miyamoto A 2005 Changes in follicular vascularity during the first follicular wave in lactating cows. *Journal of Reproduction and Development* **52** 273-280.

Armstrong DG, Gong JG & Webb R 2003 Interactions between nutrition and ovarian activity in cattle: physiological, cellular and molecular mechanisms. *Reproduction Supplement* **61** 403-414.

Beam SW & Butler WR 1997 Energy balance and ovarian follicle development prior to the first ovulation postpartum in dairy cows receiving three levels of dietary fat. *Biology of Reproduction* **56** 133-142.

Beam SW & Butler WR 1998 Energy balance, metabolic hormones, and early postpartum follicular development in dairy cows fed prilled lipid. *Journal of Dairy Science* **81** 121-131.

Beam SW & Butler WR 1999 Effects of energy balance on follicular development and first ovulation in postpartum dairy cows. *Journal of Reproduction and Fertility Supplement* **54** 411-424.

Butler ST, Pelton SH & Butler WR 2004 Insulin increases 17 beta-estradiol production by the dominant follicle of the first postpartum follicle wave in dairy cows. *Reproduction* **127** 537-545.

Canfield RW & Butler WR 1990 Energy balance and pulsatile LH secretion in early postpartum dairy cattle. *Domestic Animal Endocrinology* **7** 323-330.

Darwash AO, Lamming GE & Woolliams JA 1997 The phenotypic association between the interval to post-partum ovulation and traditional measures of fertility in dairy cattle. *Animal Science* **65** 9-16.

Daughaday WH, Mariz IK & Blethen SL 1980 Inhibition of access of bound somatomedin to membrane receptor and immunobinding sites: a comparison of radioreceptor and radioimmunoassay of somatomedin in native and acid-ethanol-extracted serum. *Journal of Clinical Endocrinology and Metabolism* **51** 781-788.

- 1 **Ferguson JD, Galligan DT & Thomsen N** 1994 Principal descriptors of body condition score in Holstein
2 cows. *Journal of Dairy Science* **77** 2695-2703.
- 3 **Fortune JE, Rivera GM & Yang MY** 2004 Follicular development: the role of the follicular
4 microenvironment in selection of the dominant follicle. *Animal Reproduction Science* **82-83** 109-126.
- 5 **Ginther OJ, Beg MA, Donadeu FX & Bergfelt DR** 2003 Mechanism of follicle deviation in monovular
6 farm species. *Animal Reproduction Science* **78** 239-257.
- 7 **Glister C, Tannetta DS, Groome NP & Knight PG** 2001 Interactions between follicle-stimulating
8 hormone and growth factors in modulating secretion of steroids and inhibin-related peptides by
9 nonluteinized bovine granulosa cells. *Biology of Reproduction* **65** 1020-1028.
- 10 **Gong JG, Lee WJ, Garnsworthy PC & Webb R** 2002 Effect of dietary-induced increases in circulating
11 insulin concentrations during the early postpartum period on reproductive function in dairy cows.
12 *Reproduction* **123** 419-427.
- 13 **Gutierrez CG, Campbell BK & Webb R** 1997 Development of a long-term bovine granulosa cell culture
14 system: induction and maintenance of estradiol production, response to follicle-stimulating hormone,
15 and morphological characteristics. *Biology of Reproduction* **56** 608-616.
- 16 **Hayashi KG, Acosta TJ, Tetsuka M, Berisha B, Matsui M, Schams D, Ohtani M & Miyamoto A** 2003
17 Involvement of angiopoietin-tie system in bovine follicular development and atresia: messenger RNA
18 expression in theca interna and effect on steroid secretion. *Biology of Reproduction* **69** 2078-2084.
- 19 **Jiang JY, Macchiarelli G, Tsang BK & Sato E** 2003 Capillary angiogenesis and degeneration in bovine
20 ovarian antral follicles. *Reproduction* **125** 211-223.
- 21 **Jolly PD, McDougall S, Fitzpatrick LA, Macmillan KL & Entwistle KW** 1995 Physiological effects of
22 undernutrition on postpartum anoestrus in cows. *Journal of Reproduction and Fertility Supplement* **49**
23 477-492.
- 24 **Kawashima C, Kaneko E, Amaya Montoya C, Matsui M, Yamagishi N, Matsunaga N, Ishii M, Kida**
25 **K, Miyake Y-I & Miyamoto A** (2006) Relationship between the first ovulation within three weeks
26 postpartum and subsequent ovarian cycles and fertility in high producing dairy cows. *Journal of*
27 *Reproduction and Development* **52** 479-486.

- 1 **Kobayashi Y, Boyd CK, Bracken CJ, Lamberson WR, Keisler DH & Lucy MC** 1999 Reduced growth
2 hormone receptor (GHR) messenger ribonucleic acid in liver of periparturient cattle is caused by a
3 specific down-regulation of GHR 1A that is associated with decreased insulin-like growth factor I.
4 *Endocrinology* **140** 3947-3954.
- 5 **Lamming GE & Bulman DC** 1976 The use of milk progesterone radioimmunoassay in the diagnosis and
6 treatment of subfertility in dairy cows. *British Veterinary Journal* **132** 507-517.
- 7 **Lamming GE, Wathes DC & Peters AR** 1981 Endocrine patterns of the post-partum cow. *Journal of*
8 *Reproduction and Fertility Supplement* **30** 155-170.
- 9 **Lucy MC, Beck J, Staples CR, Head HH, De La Sota RL & Thatcher WW** 1992a Follicular dynamics,
10 plasma metabolites, hormones and insulin-like growth factor I (IGF-I) in lactating cows with positive or
11 negative energy balance during the preovulatory period. *Reproduction Nutrition and Development* **32**
12 331-341.
- 13 **Lucy MC, Staples CR, Thatcher WW, Erickson PS, Cleale RM, Firkins JL, Clark JH, Murphy MR**
14 **& Brodie BO** 1992b Influence of diet composition, dry matter intake, milk production and fertility in
15 dairy cows. *Animal Production* **54** 323-331.
- 16 **McCann JP & Hansel W** 1986 Relationships between insulin and glucose metabolism and pituitary-
17 ovarian functions in fasted heifers. *Biology of Reproduction* **34** 630-641.
- 18 **Miyamoto A, Okuda K, Schweigert FJ & Schams D** 1992 Effects of basic fibroblast growth factor,
19 transforming growth factor-beta and nerve growth factor on the secretory function of the bovine corpus
20 luteum in vitro. *Journal of Endocrinology* **135** 103-114.
- 21 **Morimoto S, Fernandez-Mejia C, Romerro-Navarro G, Morales-Peza N & Diaz-Sanchez V** 2001
22 Insulin gene expression pattern in rat pancreas during the estrous cycle. *Life Sciences* **68** 2979-2985.
- 23 **Mutayoba B, Meyer H, Schams D & Schallenberger E** 1990 Development of a sensitive enzyme
24 immunoassay for LH determination in bovine plasma using the streptavidin-biotin technique. *Acta*
25 *Endocrinologica (Copenhagen)* **122** 227-232.
- 26 **Pushpakumara PG, Gardner NH, Reynolds CK, Beever DE & Wathes DC** 2003 Relationships between
27 transition period diet, metabolic parameters and fertility in lactating dairy cows. *Theriogenology* **60**
28 1165-1185.

- 1 **Radcliff RP, McCormack BL, Crooker BA & Lucy MC** 2003 Plasma hormones and expression of
2 growth hormone receptor and insulin-like growth factor-I mRNA in hepatic tissue of periparturient
3 dairy cows. *Journal of Dairy Science* **86** 3920-3926.
- 4 **Roh S-G, Matsunaga N, Miyamoto A, Hidaka S & Hidari H** 1997 Competitive enzyme immunoassay for
5 bovine growth hormone. *Endocrine Journal* **44** 195-198.
- 6 **Savio JD, Boland MP, Hynes N & Roche JF** 1990a Resumption of follicular activity in the early post-
7 partum period of dairy cows. *Journal of Reproduction and Fertility* **88** 569-579.
- 8 **Savio JD, Boland MP & Roche JF** 1990b Development of dominant follicles and length of ovarian cycles
9 in post-partum dairy cows. *Journal of Reproduction and Fertility* **88** 581-591.
- 10 **Senatore EM, Butler WR & Oltenu PA** 1996 Relationship between energy balance and postpartum
11 ovarian activity and fertility in first lactation dairy cows. *Animal Science* **62** 17-23.
- 12 **Spicer LJ & Stewart RE** 1996 Interactions among basic fibroblast growth factor, epidermal growth factor,
13 insulin, and insulin-like growth factor-I (IGF-I) on cell numbers and steroidogenesis of bovine thecal
14 cells: role of IGF-I receptors. *Biology of Reproduction* **54** 255-263.
- 15 **Spicer LJ, Tucker WB & Adams GD** 1990 Insulin-like growth factor-I in dairy cows: relationships
16 among energy balance, body condition, ovarian activity, and estrous behavior. *Journal of Dairy Science*
17 **73** 929-937.
- 18 **Spicer LJ, Alpizar E & Echternkamp SE** 1993 Effects of insulin, insulin-like growth factor I, and
19 gonadotropins on bovine granulosa cell proliferation, progesterone production, estradiol production,
20 and(or) insulin-like growth factor I production in vitro. *Journal of Animal Science* **71** 1232-1241.
- 21 **Staples CR, Thatcher WW & Clark JH** 1990 Relationship between ovarian activity and energy status
22 during the early postpartum period of high producing dairy cows. *Journal of Dairy Science* **73** 938-947.
- 23 **Stevenson JS & Britt JH** 1979 Relationships among luteinizing hormone, estradiol, progesterone,
24 glucocorticoids, milk yield, body weight and postpartum ovarian activity in Holstein cows. *Journal of*
25 *Animal Science* **48** 570-577.
- 26 **Watanabe H, Miyamoto A & Fukui Y** 1997 A competitive enzyme immunoassay for follicle-stimulating
27 hormone in ovine plasma using biotin-streptavidin amplification. *Reproduction, Fertility and*
28 *Development* **9** 597-601.

- 1 **Yamada O, Abe M, Takehana K, Iwasa K, Hiraga T & Hratsuka T** 1994 Microvasculature of mature
- 2 bovine follicles and its changes with ovulation. *Journal of Reproduction and Development* **40** 307-315.

1

2

Table 1. The metabolic-endocrine factors between before and after E2 peak period in ovulatory cows

	Relative to E2 peak		Significance of differences
	Before	After	
Insulin (ng/ml)	0.29±0.03	0.47±0.08	P=0.06
GH (ng/ml)	11.7±1.5	9.6±0.8	NS
IGF-1 (ng/ml)	98.2±21.5	61.1±13.7	P<0.05

Values are mean±sem

NS; not significant.

Data were divided into two phases which were before E2 peak (-10 to -1 days relative to E2 peak) and after E2 peak (0 to 10 days relative to E2 peak)

1 Figure Legends

2

3 **Figure 1** Metabolites, metabolic hormones and BCS from 4 wk prepartum to 3 wk postpartum for cows
4 that developed either ovulatory (n=13) or anovulatory (n=9) dominant follicle during the first follicular
5 wave postpartum (mean±sem, solid; ovulation, open; anovulation). The first sampling week after
6 parturition was regarded as 0 week. The serum glucose concentrations in the anovulatory cows were lower
7 than that in the ovulatory cows during the postpartum period (P<0.001). BCS in ovulatory cows tended to
8 be higher than in anovulatory cows during the postpartum period (P=0.08). The plasma concentration of
9 GH in anovulatory cows was higher than that in ovulatory cows during the peripartum period (prepartum;
10 P<0.01, postpartum; P<0.0001). On the other hand, the plasma concentration of IGF-1 in ovulatory cows
11 was higher than that in anovulatory cows during the peripartum periods (prepartum; P<0.05, postpartum;
12 P<0.01).The concentrations of NEFA, T-cho, AST and insulin were similar between the ovulatory and the
13 anovulatory cows during the period of study.

14

15 **Figure 2** Representative images of dominant follicles during the first follicular wave postpartum of
16 ovulatory and anovulatory cow. Red colour represents the blood flow toward the transducer, and blue
17 indicates blood flow away from the transducer. The colour gain of the flow mode was set to detect
18 movement of at least 2 mm/sec Scale bar represents 5 mm. The white dotted lines which delineate circles
19 indicate the frame of the follicle and the green dotted lines which delineate circles indicate the frame of the
20 corpus luteum.

21

22 **Figure 3** The function and morphology of the dominant follicle during the first follicular wave postpartum
23 and metabolic status, which was extracted data from Fig. 1, either ovulatory (n=13) or anovulatory (n=9)
24 cows at 1, 1.5 and 2 weeks postpartum (mean±sem, solid; ovulation, open; anovulation). ** indicates
25 differences of P<0.01 between ovulatory and anovulatory cows at 1.5 weeks postpartum. a and b indicates
26 differences of P<0.05 within ovulatory cows. The diameter of the dominant follicle was similar growth in
27 both groups (time effect; P<0.01) and did not differ between the ovulatory and the anovulatory cows during
28 the period of study. The plasma concentrations of E2 in ovulatory cows were higher than that in

1 anovulatory cows at 1.5 weeks postpartum ($P<0.05$), however there was no significant differences between
2 ovulatory and anovulatory at 2 weeks postpartum because of decrease of preovulatory E2 concentrations. In
3 ovulatory cows, plasma GH concentrations were lower ($P<0.01$) and IGF-1 concentrations were higher
4 ($P<0.05$) compared to anovulatory cows, however the plasma insulin concentrations did not differ between
5 the ovulatory and anovulatory cows during this period.


6

7 **Figure 4** Daily representations of plasma concentrations of insulin, GH, IGF-1 and P4 in 9 out of 22 cows.
8 We obtained blood samples 4 times/day for detection (presence or absence) of LH surge, 2 time/day for
9 detection of E2 peak and 1 time/day for profile of P4 and metabolic hormones. E2 concentrations are
10 shown at 6 h intervals from 10 d before to 10 d after E2 peak in ovulated ($n=5$) and from 10 d to 20 d
11 postpartum in anovulated cows ($n=4$) (mean \pm sem). Insulin, GH, IGF-1 and P4 concentrations are indicated
12 by squares and E2 concentrations are indicated by circles; solid in ovulated cows (■, ●) and open in
13 anovulated cows (□, ○). Dotted lines in both of graphs indicate time of E2 peak in ovulated cows, and the
14 orange vertical bar of the graph for ovulated cows indicates the range of LH surge (21 ± 7 h after E2 peak).

15


16 **Figure 5** The summary of the results as a schematic representation of the growth and maturation of the
17 dominant follicle at the first follicular wave postpartum. High IGF-1 levels may stimulate the follicular cell
18 function such as the steroidogenesis and the proliferation to appear healthy dominant follicle, and after that,
19 the levels decreases because of negative energy balance. Subsequently, E2 was more secreted.
20 Contrastively, the increase in insulin level together with E2 peak may enhance the maturation of the
21 dominant follicle. Consequently, E2 enhanced by IGF-1 induce LH surge and the dominant follicle reaches
22 ovulation.

1 Figure 1




1 Figure 2

Ovulatory cow



Anovulatory cow



7 days pp


14 days pp

17 days pp


21 days pp

5mm
┆

1 Figure 3



1 Figure 4



1 Figure 5

2

3

