

Research Briefs

Title

*Babesia bovis*: Effects of Cysteine Protease Inhibitors on *In Vitro* Growth

Authors

Kazuhiro Okubo, Naoaki Yokoyama,\* Yadav Govind,<sup>†</sup> Andy Alhassan, and Ikuo Igarashi

Address

National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary  
Medicine, Inada-cho, Obihiro, Hokkaido 080-8555, Japan

\*Corresponding Author

Naoaki Yokoyama: National Research Center for Protozoan Diseases, Obihiro University of  
Agriculture and Veterinary Medicine, Inada-cho, Obihiro, Hokkaido 080-8555, Japan  
TEL.: +81-155-49-5649; FAX: +81-155-49-5643; E-mail: yokoyama@obihiro.ac.jp

<sup>†</sup>Present address

Department of Epidemiology and Preventive Medicine, College of Veterinary Science, Mathura UP. Pt.  
Deendayal Upadhaya Veterinary University, Mathura-281001(U.P), India

Abstract

In the present study, we examined the effects of four kinds of cysteine protease inhibitors (E64, E64d, leupeptin, and ALLN) on the *in vitro* asexual growth of *Babesia bovis*. Of these, only the lipophilic inhibitors, E64d and ALLN, were found to effectively inhibit the growth of *B. bovis*. In further experiments, E64d, but not ALLN, significantly suppressed the parasite's invasion of host erythrocytes, while both chemicals, especially ALLN, inhibited the parasite's replication within the infected erythrocytes. These data suggested the presence of cysteine protease(s) derived from *B. bovis*, in which the protease(s) would play important roles in the erythrocyte invasion and/or replication processes of the parasite.

Key words: *Babesia bovis*, cysteine protease, inhibitor, invasion, replication.

*Babesia bovis*, an obligatory intraerythrocytic parasite of the phylum Apicomplexa, is a major causative agent of bovine babesiosis, which causes severe clinical symptoms, such as fever, anemia, and cerebral dysfunctions, in cattle due to their asexual growth (Homer *et al.*, 2000). The disease often results in great economic losses in the livestock industry worldwide (Brown and Palmer, 1999), and effective strategies are desired for eradicating babesiosis.

Cysteine proteases are known to play vital roles in the growth of several protozoan parasites. In *Trypanosoma* spp., for example, the cysteine protease-specific inhibitors impair their host cell invasion and arrest the intracellular development (Meirelles *et al.*, 1992) or kill the cultured blood stream forms of the parasites (Troeberg *et al.*, 1999). In *Plasmodium* spp., several protozoan cysteine proteases are involved in the erythrocyte invasion (Greenbaum *et al.*, 2002), the degradations of hemoglobin and erythrocyte cytoskeletal proteins (Sijwali and Rosenthal, 2004; Takakuwa, 2001), and the final erythrocyte rupture for their egression (Wickham *et al.*, 2003). Cysteine protease inhibitors have been studied for the development of new chemotherapeutic measures for trypanosomiasis (Engel *et al.*, 1998), malaria (Olson *et al.*, 1999), schistosomiasis (Wasilewski *et al.*, 1996), and leishmaniasis (Das *et al.*, 2001). However, no cysteine protease has been reported in *Babesia* parasites, except for “cys1,” which is a cathepsin-like cysteine protease found in *Babesia (Theileria) equi* (Eakin *et al.*, 1990; Holman *et al.*, 2002; Mehlhorn and Schein, 1998).

In the present study, we examined the inhibitory effects of four commercial cysteine protease inhibitors, E64 (*trans*-Epoxy-succinyl-L-leucylamido(4-guanidino)butane), E64d ((2S,3S)-*trans*-Epoxy-succinyl-L-leucylamido-3-methylbutane ethyl ester), leupeptin, and ALLN (*N*-acetyl-leucine-leucine-norleucinal), on the *in vitro* asexual growth of cultured *B. bovis*. These inhibitors are known to inhibit thiol proteases, such as calpain and cathepsin, but only leupeptin can also inhibit serine proteases (Carole and Wang, 1998). Furthermore, E64d and ALLN have membrane permeability, while E64 and leupeptin do not (Holman *et al.*, 2002). These four inhibitors have been widely used in the research of the cysteine proteases of many protozoa, including a *Plasmodium* parasite (Dahl and Rosenthal, 2005), which is closely related to the *Babesia* parasite.

The Texas strain of *B. bovis* was cultured in purified bovine erythrocytes (RBCs) using a

serum-free GIT medium (Wako Pure Chemical Industries, Ltd., Osaka, Japan) as described previously (Bork *et al.*, 2005). The growth-inhibition test of four cysteine protease inhibitors followed previously described methods for measuring the drug activity (Okubo *et al.*, 2006a; Bork *et al.*, 2003) with some modifications. Briefly, infected RBCs were diluted with non-infected RBCs to obtain 0.5% parasitemia. Fifty  $\mu$ l of the infected RBC mixture was subsequently suspended in 450  $\mu$ l of a culture medium supplemented with 1 (or 20) - 220  $\mu$ M of E64, E64d, leupeptin, or ALLN. The suspension was placed in a 48-well culture plate (Nunc A/S, Roskilde, Denmark) and then incubated in a humidified multigas water-jacketed incubator at 37°C in 5% CO<sub>2</sub>, 5% O<sub>2</sub>, and 90% N<sub>2</sub> for 3 days. In parallel, normal cultures, which were added with the same final volume of DMSO (solvent) instead of each chemical, were prepared as the control. The culture medium was replaced daily with 450  $\mu$ l of the fresh medium containing the indicated concentration of each chemical.

Out of the four tested cysteine protease inhibitors, only the lipophilic inhibitors, E64d and ALLN, significantly affected the asexual growth of *B. bovis* at the concentration of 50  $\mu$ M (Fig. 1). On the other hand, the hydrophilic inhibitors, E64 and leupeptin, did not affect the growth of the parasite in the range of 20 - 200  $\mu$ M (Fig. 1; data not shown). The IC<sub>50</sub> values of E64d and ALLN were calculated to be 29.9 and 25.1  $\mu$ M, respectively, on the basis of the inhibitory rates (%) of E64d and ALLN at various concentrations within 1 - 220  $\mu$ M (Table 1). These data suggest that *B. bovis* has cysteine protease(s). On the other hand, E64 and leupeptin did not show any effects on the asexual growth of *B. bovis* in the present study. One possible explanation of the negative results is that the target cysteine protease(s) might exist in the parasite's body; therefore, the membrane non-permeable agents could not reach it. Another possible explanation is that the E64 and leupeptin inhibited the target cysteine protease(s) but the protease(s) did not play an essential role in the asexual growth of the parasite. In *P. falciparum*, for example, targeted disruption of the cysteine protease, falcipain 1, did not influence their asexual growth (Saliha *et al.*, 2004).

Next, we investigated the effects of E64d and ALLN on the erythrocyte invasion of *B. bovis* by an *in vitro* invasion test using high-voltage pulsing, as described previously (Okubo *et al.*, 2006a; Franssen *et al.*, 2003), with some modifications. Briefly, after *B. bovis*-infected RBCs were suspended

1 in an equal volume of a GIT medium, the mixture of 400 µl was subjected to five intermittent (10 sec,  
2 4°C) high-voltage pulses (1.5 kV, 400 Ω, 25 µF) in a BioRad Gene Pulser II (Hercules, CA, USA) with  
3 a 0.2 cm pulser cuvette (BioRad) to rupture all of the infected and non-infected RBCs exclusively and  
4 prepare extraerythrocytic (free) parasites. The samples were then suspended in a GIT medium  
5 supplemented with E64d or ALLN at the indicated concentrations (20, 50, and 100 µM) or with the  
6 same final concentration of DMSO for the control. After a low centrifugation at 700 x g for 3 min, the  
7 pellet was resuspended in the same medium supplemented with E64d, ALLN, or DMSO and then  
8 transferred into a 96-well culture plate (Nunc A/S) with non-infected RBCs at a 10% packed cell  
9 volume. After the incubation at 37°C for 1 h, the number of infected RBCs was counted out of a total  
10 5,000 RBCs in the Giemsa-stained smears, and the invasion efficiency was calculated as the  
11 percentage of parasitemia in the culture with E64d or ALLN to that with a DMSO medium control  
12 (100%).

13 As a result, the erythrocyte invasion activity of the parasite was significantly inhibited in the  
14 presence of 100 and 50 µM E64d ( $P = 0.010$  and  $0.045$ , respectively) (Fig. 2A). In contrast, none of the  
15 concentrations (100, 50, and 20 µM) of ALLN showed any significant inhibitory effects relative to the  
16 value of the control (Fig. 2B). Although the other two inhibitors (E64 and leupeptin) were also tested  
17 on the invasion and following replication tests, no inhibitory effects on these activities were noted (data  
18 not shown). This finding suggests the presence of an anonymous but functional babesial cysteine  
19 protease(s) that appears to be essential for the parasite's invasion to host RBCs. In contrast, the lack of  
20 effect of ALLN might be explained by the lack of activity of this inhibitor over the particular cysteine  
21 protease(s). In *P. falciparum*, it is known that a parasite's cysteine protease, falcipain 1, plays a specific  
22 role in host cell invasion and is also inhibited by E64d (Greenbaum *et al.*, 2002; Dahl and Rosenthal,  
23 2005). Our data will be useful for elucidating the molecular mechanism of the erythrocyte invasion of  
24 *Babesia* parasites in future.

25 Finally, the effects of E64d and ALLN on the intraerythrocytic replication of *B. bovis* were also  
26 evaluated using the high-voltage pulsing method described previously (Okubo *et al.*, 2006b). After  
27 high-voltage pulsing, free parasites were incubated with normal RBCs at 37°C. When the parasites

1 were incubated for 30 min, almost all parasites invaded the RBCs and were observed to form rings on  
2 the stained smears (data not shown, Okubo *et al.*, 2006b). At that time, the infected RBCs were washed  
3 with the GIT medium and then incubated with the indicated concentrations (20, 50, and 100  $\mu$ M) of  
4 E64d or ALLN or with the same final concentration of DMSO for the control, at 37°C for 5 h. After the  
5 incubation, Giemsa-stained smears were prepared, and the replication activity was calculated as the  
6 ratio of the number of divided parasite-containing RBCs to the entire population of infected RBCs  
7 (100%), among which more than 200 of the infected RBCs were monitored. The divided  
8 parasite-containing RBCs are infected RBCs with more than two replicated parasites before their  
9 egression (data not shown).

10 As shown in Figure 3, supplements with 100 and 50  $\mu$ M E64d had stronger inhibitory effects on  
11 the parasite's replication than the control culture ( $P = 0.0019$  and  $0.018$ , respectively) (Fig. 3A).  
12 Moreover, 100, 50, and 20  $\mu$ M ALLN had more significant inhibitory effects on the replication activity  
13 ( $P = 0.00026$ ,  $0.00029$ , and  $0.00038$ , respectively) (Fig. 3B) than E64d. In *P. falciparum*, the parasite's  
14 cysteine proteases have been reported to play a vital role in host hemoglobin degradation (Sijwali and  
15 Rosenthal, 2004). Both E64d and ALLN were reported to block the intracellular processing of two  
16 parasite's cysteine proteases, falcipain-2 and -3 (Dahl and Rosenthal, 2005), which function to degrade  
17 the host hemoglobin in the intracellular stage (Sijwali *et al.*, 2001; Shenai *et al.*, 2000). Accordingly, it  
18 is also possible that the anonymous babesial cysteine protease could digest the host hemoglobin for  
19 their metabolism even if the phenomenon has not been observed in *Babesia* parasites.

20 Consequently, our data strongly suggested the presence of cysteine protease(s) derived from *B.*  
21 *bovis*, in which the protease(s) would play important roles in the erythrocyte invasion and/or  
22 replication processes of *B. bovis*. However, since there is no report about the identification of *B. bovis*  
23 cysteine protease(s) related to their asexual growth or even the homologous proteins, the exact role of  
24 babesial cysteine protease is still not clear. Progress of the genome project is expected to lead to the  
25 elucidation of the *B. bovis* cysteine protease. Further research will be necessary to elucidate the  
26 molecular mechanism of the asexual growth cycle of *Babesia* parasites as well as for developing novel  
27 anti-babesial drugs in the future.

## Acknowledgments

This study was supported by Grants-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (JSPS), the Program for the Promotion of Basic Research Activities for Innovative Biosciences (PROBRAIN), The 21<sup>st</sup> Century COE Program (A-1), Ministry of Education, Culture, Sports, Science, and Technology, and the Japan International Cooperation Agency (JICA), Japan. A. A. was supported by a research fellowship from JSPS.

## References

- BORK, S., OKAMURA, M., MATSUO, T., KUMAR, S., YOKOYAMA, N., and IGARASHI, I. 2005. Host serum modifies the drug susceptibility of *Babesia bovis* *in vitro*. *Parasitology* 130, 489-492.
- BORK, S., YOKOYAMA, N., MATSUO, T., CLAVERIA, F. G., FUJISAKI, K., and IGARASHI, I. 2003. Growth inhibitory effect of triclosan on equine and bovine *Babesia* parasites. *American Journal of Tropical Medicine and Hygiene* 68, 334-340.
- BROWN, W. C., and PALMER, G. H. 1999. Designing blood-stage vaccines against *Babesia bovis* and *B. bigemina*. *Parasitology Today* 15, 275-281.
- CAROLE, L. M., and Kuan, W. 1998. Effect of thiol protease inhibitors on myoblast fusion and myofibril assembly *in vitro*. *Cell Motility and the Cytoskeleton* 40, 354-367.
- DAHL, E. L., and ROSENTHAL, P. J. 2005. Biosynthesis, localization, and processing of falcipain cysteine proteases of *Plasmodium falciparum*. *Molecular and Biochemical Parasitology* 139, 205-212.
- DAS, L., DATTA, N., BANDYOPADHYAY, S., and DAS, P. K. 2001. Successful therapy of lethal murine visceral leishmaniasis with cystatin involves up-regulation of nitric oxide and a favorable T cell response. *Journal of Immunology* 166, 4020-4028.
- EAKIN, A. E., BOUVIER, J., SAKANARI, J. A., CRAIK, C. S., and MCKERROW, J. H. 1990. Amplification and sequencing of genomic DNA fragments encoding cysteine proteases from protozoan parasites. *Molecular and Biochemical Parasitology* 39, 1-8.
- ENGEL, J. C., DOYLE, P. S., HSIEH, I., and MCKERROW, J. H. 1998. Cysteine protease inhibitors cure an experimental *Trypanosoma cruzi* infection. *Journal of Experimental Medicine* 188, 725-734.
- FRANSSEN, F. F., GAFFAR, F. R., YATSUDA, A. P., and DE VRIES, E. 2003. Characterization of

erythrocyte invasion by *Babesia bovis* merozoites efficiently released from their host cell after high-voltage pulsing. *Microbes and Infection* 5, 365-372.

GREENBAUM, D. C., BARUCH, A., GRAINGER, M., BOZDECH, Z., MEDZIHRADESKY, K. F., ENGEL, J., DERISI, J., HOLDER, A. A., and BOGYO, M. 2002. A role for the protease falcipain 1 in host cell invasion by the human malaria parasite. *Science* 298, 2002-2006.

HOLMAN, P. J., HSIEH, M. M., NIX, J. L., BENDELE, K. G., WAGNER, G. G., and BALL, J. M. 2002. A cathepsin L-like cysteine protease is conserved among *Babesia equi* isolates. *Molecular and Biochemical Parasitology* 119, 295-300.

HOMER, M. J., AGUILAR-DELFIN, I., TELFORD, S. R., 3RD, KRAUSE, P. J., and PERSING, D. H. 2000. Babesiosis. *Clinical Microbiology Reviews* 13, 451-469.

MEHLHORN, H., and SCHEIN, E. 1998. Redescription of *Babesia equi* Laveran, 1901 as *Theileria equi* Mehlhorn, Schein 1998. *Parasitology Research* 84, 467-475.

MEIRELLES, M. N., JULIANO, L., CARMONA, E., SILVA, S. G., COSTA, E. M., MURTA, A. C., and SCHARFSTEIN, J. 1992. Inhibitors of the major cysteinyl proteinase (GP57/51) impair host cell invasion and arrest the intracellular development of *Trypanosoma cruzi* *in vitro*. *Molecular and Biochemical Parasitology* 52, 175-184.

OKUBO, K., WILAWAN, P., BORK, S., OKAMURA, M., YOKOYAMA, N., and IGARASHI, I. 2006a. Calcium-ions are involved in erythrocyte invasion by equine *Babesia* parasites. *Parasitology* 133, 289-294.

OKUBO, K., YOKOYAMA, N., TAKABATAKE, N., and IGARASHI, I. 2006b. Amount of cholesterol in host membrane affects erythrocyte invasion and replication by *Babesia bovis*. *Parasitology* 134, 1-6.

OLSON, J. E., LEE, G. K., SEMENOV, A., and ROSENTHAL, P. J. 1999. Antimalarial effects in

mice of orally administered peptidyl cysteine protease inhibitors. *Bioorganic and Medicinal Chemistry* 7, 633-638.

SALIHA, E., BEATA, C., DORON, C. G., MATTHEW, B., and KIM, C. W. 2004. Targeted disruption of *Plasmodium falciparum* cysteine protease, falcipain 1, reduces oocyst production, not erythrocyte stage growth. *Molecular Microbiology* 53, 243-250.

SHENAI, B. R., SIJWALI, P. S., SINGH, A., and ROSENTHAL, P. J. 2000. Characterization of native and recombinant falcipain-2, a principal trophozoite cysteine protease and essential hemoglobinase of *Plasmodium falciparum*. *The Journal of Biological Chemistry* 275, 29000-29010.

SIJWALI, P. S., and ROSENTHAL, P. J. 2004. Gene disruption confirms a critical role for the cysteine protease falcipain-2 in hemoglobin hydrolysis by *Plasmodium falciparum*. *Proceedings of the National Academy of Science of the United States of America* 101, 4384-4389.

SIJWALI, P. S., SHENAI, B. R., GUT, J., SINGH, A., and ROSENTHAL, P. J. 2001. Expression and characterization of the *Plasmodium falciparum* hemoglobinase falcipain-3. *The Biochemical Journal* 360, 481-489.

TAKAKUWA, Y. 2001. Regulation of red cell membrane protein interactions: Implications for red cell function. *Current Opinion in Hematology* 8, 80-84.

TROEBERG, L., MORTY, R. E., PIKE, R. N., LONSDALE-ECCLES, J. D., PALMER, J. T., MCKERROW, J. H., and COETZER, T. H. 1999. Cysteine proteinase inhibitors kill cultured bloodstream forms of *Trypanosoma brucei brucei*. *Experimental Parasitology* 91, 349-355.

WASILEWSKI, M. M., LIM, K. C., PHILLIPS, J., and MCKERROW, J. H. 1996. Cysteine protease inhibitors block schistosome hemoglobin degradation *in vitro* and decrease worm burden and egg production *in vivo*. *Molecular and Biochemical Parasitology* 81, 179-189.

WICKHAM, M. E., CULVENOR, J. G., and COWMAN, A. F. 2003. Selective inhibition of a


- 1 two-step egress of malaria parasites from the host erythrocyte. The Journal of Biological Chemistry
- 2 278, 37658-37663.
- 3

## Figure Legends



**Fig. 1.** Effects of four cysteine protease inhibitors on the *in vitro* growth of *B. bovis*. Each value represents the mean  $\pm$  standard deviation (SD) in 3 wells for each 50  $\mu$ M chemical (E64, E64d, leupeptin, or ALLN) in 3 separate experiments. The asterisks indicate significant differences (\* $P$  < 0.05, \*\* $P$  < 0.005) between the inhibitor- and DMSO control-treated parasitemia analyzed using an independent Student's t-test.

**Fig. 2.** Effects of E64d (A) and ALLN (B) on the erythrocyte invasion of *B. bovis*. Relative values are expressed as the percentage of the parasitemia in the culture with E64d or ALLN to that in the DMSO medium control (Con: 100%) in an *in vitro* invasion test. Each value represents the mean  $\pm$  SD in 3 wells for each concentration of chemicals in 3 separate experiments. The asterisks indicate significant differences (\* $P$  < 0.05) on the values between the chemical- and DMSO control-treated cultures analyzed using an independent Student's t-test.



**Fig. 3.** Effects of E64d (A) and ALLN (B) on the intraerythrocytic replication of *B. bovis*. Relative values are expressed as the rates of divided parasite-containing RBCs to all infected RBCs. Each value represents the mean  $\pm$  SD in 3 wells for each concentration of chemicals in 3 separate experiments. The asterisks indicate significant differences (\* $P$  < 0.05, \*\* $P$  < 0.005) on the values between the chemical- and DMSO control-treated cultures analyzed using an independent Student's t-test.



**Fig. 1. Okubo *et al.***



**Fig. 2. Okubo *et al.***



**Fig. 3. Okubo *et al.***